

Modular Symmetry and Fractional Charges in $\mathcal{N} = 2$ Supersymmetric Yang-Mills and the Quantum Hall Effect

Brian P. Dolan [†]

[†] *Department of Mathematical Physics, National University of Ireland, Maynooth, Ireland*
and
School of Theoretical Physics, Dublin Institute for Advanced Studies, 10, Burlington Rd., Dublin, Ireland

E-mail: bdolan@thphys.nuim.ie

URL: <http://www.thphys.nuim.ie/staff/bdolan/>

Received ???, in final form ???; Published online ???

Original article is available at [http://www.emis.de/journals/SIGMA/2006/Paper?/?/](http://www.emis.de/journals/SIGMA/2006/Paper?/)

Abstract. The parallel rôles of modular symmetry in $\mathcal{N} = 2$ supersymmetric Yang-Mills and in the quantum Hall effect are reviewed. In supersymmetric Yang-Mills theories modular symmetry emerges as a version of Dirac's electric – magnetic duality. It has significant consequences for the vacuum structure of these theories, leading to a fractal vacuum which has an infinite hierarchy of related phases. In the case of $\mathcal{N} = 2$ supersymmetric Yang-Mills in 3+1 dimensions, scaling functions can be defined which are modular forms of a sub-group of the full modular group and which interpolate between vacua. Infra-red fixed points at strong coupling correspond to θ -vacua with θ a rational number that, in the case of pure SUSY Yang-Mills, has odd denominator. There is a mass gap for electrically charged particles which can carry fractional electric charge. A similar structure applies to the 2+1 dimensional quantum Hall effect where the hierarchy of Hall plateaux can be understood in terms of an action of the modular group and the stability of Hall plateaux is due to the fact that odd denominator Hall conductivities are attractive infra-red fixed points. There is a mass gap for electrically charged excitations which, in the case of the fractional quantum Hall effect, carry fractional electric charge.

Key words: duality; modular symmetry; supersymmetry; quantum Hall effect.

2000 Mathematics Subject Classification: 11F11; 81R05; 81T60; 81V70

1 Introduction

The rôle of the modular group as a duality symmetry in physics has gained increasing prominence in recent years, not only through string theory considerations and in supersymmetric gauge theories but also in 2+1 dimensional $U(1)$ gauge theories. In this last case there is considerable contact with experiment via the quantum Hall effect, where modular symmetry relates the integer and fractional quantum Hall effects. This article is a review of the current understanding of modular symmetry in the quantum Hall effect and the remarkable similarities with 3+1 dimensional $\mathcal{N} = 2$ supersymmetric Yang-Mills theory. It is an extended version of the mini-review [1].

Mathematically the connection between $\mathcal{N} = 2$ supersymmetric $SU(2)$ Yang-Mills and the quantum Hall effect lies in the constraints that modular symmetry places on the scaling flow of the two theories, as will be elucidated below. The quantum Hall effect is of course a real

physical system with imperfections and impurities and as such modular symmetry can only be approximate in the phenomena, nevertheless there is strong experimental support for the relevance of modular symmetry in the experimental data.

The significance of the modular group, as a generalisation of electric-magnetic duality, was emphasised by Shapere and Wilczek [2], and its full power was realised by Seiberg and Witten in supersymmetric Yang-Mills, motivated in part by an idea of Montonen and Olive [3]. In the case of $\mathcal{N} = 2$ SUSY Yang-Mills Seiberg and Witten showed, in their seminal papers in 1994 [4, 5], that a remnant of the full modular group survives in the low energy physics.

The earliest appearance of modular symmetry in the condensed matter literature was in the work of Cardy and Rabinovici, [6, 7] where a coupled clock model was analysed, interestingly with a view to gaining insight into quantum chromodynamics. A suggested link between phase diagrams for one-dimensional clock models and the quantum Hall effect was made in [8]. An action of the modular group leading to a fractal structure was found in applying dissipative quantum mechanics to the Hofstadter model, [9, 10] and fractal structures have also emerged in other models of the quantum Hall effect [11]. The first mention of the modular group in relation to the quantum Hall effect appears to have been by Wilczek and Shapere in [2], but these authors focused on a particular subgroup which has fixed points that are not observed in the experimental data on the quantum Hall effect. A more detailed analysis was undertaken by Lütken and Ross [12, 13] and the correct sub-group was finally identified unambiguously in [13, 14], at least for spin-split quantum Hall samples. Almost at the same time as Lütken and Ross' paper Kivelson, Lee and Zhang derived their "Law of Corresponding States" [15], relating different quantum Hall plateaux in spin-split samples. Although they did not mention modular symmetry in their paper, their map is in fact the group $\Gamma_0(2)$ described below and discussed in [13] and [14].

In §2 duality in electromagnetism is reviewed and the Dirac-Schwinger-Zwanziger quantisation condition and the Witten effect are described. Section 3 introduces the sub-groups of the modular group that are relevant to $\mathcal{N} = 2$ SUSY $SU(2)$ Yang-Mills and scaling functions, modular forms that are regular at all the singular points in the moduli space of vacua, are discussed. In §4 the relevance of the modular group to the quantum Hall effect is described and Kivelson, Lee and Zhang's derivation of the Law of Corresponding States, for spin-split quantum Hall samples, and its relation to the modular group, is explained. Modular symmetries of spin-degenerate samples are also reviewed and predictions for hierarchical structures in bosonic systems, based on a different sub-group of the modular group to that of the QHE, is also explained. Finally §5 contains a summary and conclusions.

2 Duality in Electromagnetism

Maxwell's equations in the absence of sources

$$\begin{aligned} \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} &= 0 & \nabla \cdot \mathbf{E} &= 0 \\ \nabla \times \mathbf{B} - \frac{\partial \mathbf{E}}{\partial t} &= 0 & \nabla \cdot \mathbf{B} &= 0 \end{aligned} \tag{1}$$

(using units in which $\epsilon_0 = \mu_0 = c = 1$) are not only symmetric under the conformal group in 3+1 dimensions but also under the interchange of electric field \mathbf{E} and the magnetic field \mathbf{B} , more specifically Maxwell's equations are symmetric under the map

$$\mathbf{E} \rightarrow \mathbf{B} \quad \text{and} \quad \mathbf{B} \rightarrow -\mathbf{E}. \tag{2}$$

This symmetry is known as *duality*, for any field configuration (\mathbf{E}, \mathbf{B}) there is a dual configuration $(\mathbf{B}, -\mathbf{E})$ (a very good introduction to these ideas is [16]). Duality is a useful symmetry, *e.g.* in

rectangular wave-guide problems, using this symmetry one can immediately construct transverse magnetic modes once transverse electric modes are known.

In fact there is a larger continuous symmetry which is perhaps most easily seen by defining the complex field

$$\mathbf{F} = \mathbf{B} + i\mathbf{E} \quad (3)$$

and writing the source free Maxwell's equations as

$$\nabla \times \mathbf{F} + i\frac{\partial \mathbf{F}}{\partial t} = 0, \quad \nabla \cdot \mathbf{F} = 0. \quad (4)$$

Then the map $\mathbf{F} \rightarrow e^{i\phi}\mathbf{F}$, with ϕ a constant phase, is also a symmetry of the vacuum Maxwell equations sending

$$\mathbf{B} \rightarrow \cos \phi \mathbf{B} - \sin \phi \mathbf{E} \quad \text{and} \quad \mathbf{E} \rightarrow \cos \phi \mathbf{E} + \sin \phi \mathbf{B},$$

the particular case of $\phi = \pi/2$ giving (2) above.

When electric sources, *i.e.* a current J^μ , are included this is no longer a symmetry but, in a seminal paper [17], Dirac showed that a vestige of (2) remains provided magnetic monopoles are introduced, or more generally magnetic currents, \tilde{J}^μ . By quantising a charged particle, with electric charge Q , in a background magnetic monopole field, generated by a monopole with charge M , Dirac showed that single-valuedness of the wave-function requires that QM must be an integral multiple of Planck's constant, or

$$QM = 2\pi\hbar n \quad (5)$$

where n is an integer — the famous Dirac quantisation condition which is intimately connected with topology and the theory of fibre bundles [18].

A quick way of deriving (5) is to consider the orbital angular momentum of particle of mass m and charge Q in the presence of a magnetic monopole M generating the magnetic field

$$\mathbf{B} = \frac{M}{4\pi} \frac{\mathbf{r}}{r^3}. \quad (6)$$

The form of the Lorentz force, $m\ddot{\mathbf{r}} = Q(\dot{\mathbf{r}} \times \mathbf{B})$, implies that orbital angular momentum, $\mathbf{L} = m(\mathbf{r} \times \dot{\mathbf{r}})$, is not conserved

$$\dot{\mathbf{L}} = m(\mathbf{r} \times \ddot{\mathbf{r}}) = \mathbf{r} \times (Q(\dot{\mathbf{r}} \times \mathbf{B})) = \frac{QM}{4\pi r^3} \mathbf{r} \times (\dot{\mathbf{r}} \times \mathbf{r}) = \frac{d}{dt} \left(\frac{QM}{4\pi} \frac{\mathbf{r}}{r} \right). \quad (7)$$

However

$$\mathbf{J} := \mathbf{L} - \left(\frac{QM}{4\pi} \frac{\mathbf{r}}{r} \right), \quad (8)$$

is conserved and so we define this to be the total angular momentum of the particle plus the field. Indeed

$$\mathbf{J}^{em} = \int \mathbf{r}' \times (\mathbf{E}(\mathbf{r}') \times \mathbf{B}(\mathbf{r}')) d^3 r' = - \left(\frac{QM}{4\pi} \frac{\mathbf{r}}{r} \right) \quad (9)$$

is the angular momentum of the electromagnetic field generated by the particle and the monopole when they are separated by \mathbf{r} . Assuming that \mathbf{J}^{em} is quantised in the usual way the result of a measurement should always yield

$$J^{em} = \frac{1}{2} n \hbar \quad (10)$$

where n is an integer, leading to (5).

The Dirac quantisation condition has the fascinating consequence that, if a single magnetic monopole M exists anywhere in the universe then electric charge

$$Q = \frac{2\pi\hbar n}{M} \quad (11)$$

must be quantised as a multiple of $\frac{2\pi\hbar}{M}$. Conversely if $Q = e$ is the unit electric charge then there is a unit of magnetic charge, namely

$$m = \frac{2\pi\hbar}{e}, \quad (12)$$

and the allowed magnetic charges are integral multiples of m .

Since magnetic monopoles have never been observed, if they exist at all, they must be very heavy, much heavier than an electron,¹ and so duality is not a manifest symmetry of the physics — it is at best a map between two descriptions of the same theory. In perturbation theory, since $\alpha_e = \frac{e^2}{4\pi\hbar} \approx \frac{1}{137}$ is so small, the ‘magnetic’ fine structure constant

$$\alpha_m = \frac{m^2}{4\pi\hbar} = \frac{1}{4} \frac{1}{\alpha_e} \approx 34 \quad (13)$$

is very large and magnetic monopoles have very strong coupling to the electromagnetic field. One could not construct a perturbation theory of magnetic monopoles, but one could analyse a theory of magnetic monopoles by first performing perturbative calculations of electric charges and then using duality to map the results to the strongly coupled magnetic charges. Duality is thus potentially a very useful tool in quantum field theory since it provides a mathematical tool for studying strongly interacting theories.

There is a generalisation of the Dirac quantisation condition for particles that carry both an electric and a magnetic charge at the same time, called *dyons*. If a dyon with electric charge Q and magnetic charge M orbits a second dyon with charges Q' and M' , then Schwinger and Zwanziger showed that [20, 21]

$$QM' - Q'M = 2\pi n\hbar. \quad (14)$$

A dimensionless version of the Dirac-Schwinger-Zwanziger quantisation condition can be obtained by writing

$$Q = n_e e, \quad Q' = n'_e e, \quad M = n_m m, \quad \text{and} \quad M' = n'_m m \quad (15)$$

giving

$$n_e n'_m - n'_e n_m = n \quad (16)$$

with n_e, n'_e, n_p, n'_p and n integers.

The simple \mathbf{Z}_2 duality map (13)

$$2\alpha_e \rightarrow 2\alpha_m = \frac{1}{2\alpha_e} \quad (17)$$

can be extended to a much richer structure [2, 22, 23, 24] involving an infinite discrete group, the modular group $Sl(2, \mathbf{Z})/\mathbf{Z}_2 \cong \Gamma(1)$, by including a topological term in the 4-dimensional action,

$$\begin{aligned} S &= \int \left(-\frac{1}{4e^2} F_{\mu\nu} F^{\mu\nu} + \frac{\theta}{32\pi^2} \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma} \right) d^4x \\ &= \int \left(-\frac{1}{2e^2} F \wedge *F + \frac{\theta}{8\pi^2} F \wedge F \right), \end{aligned} \quad (18)$$

¹For a review of the current status of magnetic monopoles see [19].

with $F = \frac{1}{2}F_{\mu\nu}dx^\mu \wedge dx^\nu$ in differential form notation. Define the complex parameter

$$\tau = \frac{\theta}{2\pi} + \frac{2\pi i}{e^2}, \quad (19)$$

using units with $\hbar = 1$,² and the combinations

$$F_\pm = \frac{1}{2}(F \pm i * F) \quad \Rightarrow \quad *F_\pm = \mp i F_\pm, \quad (20)$$

then the action can be written as

$$S = \frac{i}{4\pi} \int \{ \tau(F_+ \wedge *F_+) - \bar{\tau}(F_- \wedge *F_-) \}. \quad (21)$$

In terms of τ (17) generalises to a map that we shall denote by \mathcal{S}

$$\mathcal{S} : \tau \quad \rightarrow \quad -\frac{1}{\tau}, \quad (22)$$

which reduces to (17) when $\theta = 0$. Notice that the imaginary part of τ satisfies $\Im\tau > 0$ (for stability $e^2 > 0$) and so τ parameterises the upper-half complex plane.

While it is well known that the topological term in (18) has interesting physical consequences in non-abelian gauge theories, leading to the θ -vacua in QCD [25, 26, 27, 28, 29], it is not usually considered in an abelian theory. But it can have non-trivial consequences for the partition function even for abelian theories, for example if space-time is a 4-torus $T^4 = T^2 \times \tilde{T}^2$ then the second term in (18) can be non-zero. Consider for example a field configuration in which the 2-form F is the direct sum of a monopole with first Chern class p_1 on one 2-torus T^2 and a monopole with first Chern class p_2 on the second, *i.e.*

$$F = F^{(1)} + F^{(2)} \quad (23)$$

with

$$\frac{1}{2\pi} \int_{T^2} F^{(1)} = p_1 \quad (24)$$

on the first torus and

$$\frac{1}{2\pi} \int_{\tilde{T}^2} F^{(2)} = p_2 \quad (25)$$

on the second. Then

$$\frac{1}{8\pi^2} \int_{T^4} F \wedge F = \frac{1}{4\pi^2} \int_{T^2 \times \tilde{T}^2} F^{(1)} \wedge F^{(2)} = \left(\frac{1}{2\pi} \int_{T^2} F^{(1)} \right) \left(\frac{1}{2\pi} \int_{\tilde{T}^2} F^{(2)} \right) = p_1 p_2 \quad (26)$$

is an integer and e^{iS} , and hence the partition function, is invariant under $\theta \rightarrow \theta + 2\pi$. We thus define a second operation \mathcal{T} on τ

$$\mathcal{T} : \tau \quad \rightarrow \quad \tau + 1. \quad (27)$$

Both \mathcal{S} and \mathcal{T} preserve the property that $\Im\tau > 0$ and they generate the modular group acting on the upper-half complex plane. A general element of the modular group can be represented as a 2×2 matrix of integers with unit determinant,

$$\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \quad (28)$$

² $\tau = \frac{\theta}{2\pi} + \frac{2\pi i \hbar}{e^2}$ when \hbar is included.

with $ad - bc = 1$, and

$$\gamma(\tau) = \frac{a\tau + b}{c\tau + d}. \quad (29)$$

Clearly $-\gamma$ has the same effect on τ as γ does so the modular group is $\Gamma(1) \cong Sl(2, \mathbf{Z})/\mathbf{Z}_2$. In Euclidean space a consequence of this is that the partition transforms is related to a modular form and depends on topological invariants, the Euler characteristic and the Hirzebruch signature, in a well-defined manner, [23, 24].

Magnetic monopoles *à la* Dirac are singular at the origin but, remarkably, it is possible to find classical solutions of coupled $SO(3)$ Yang-Mills-Higgs systems which are finite and smooth at the origin and reduce to monopole configurations, with $M = -\frac{4\pi\hbar}{e}$, at large r , 't Hooft-Polyakov monopoles [30, 31]. The Higgs field Φ , which is in the adjoint representation of $SO(3)$, acquires a non-zero vacuum expectation value away from the origin, $\Phi \cdot \Phi = \mathbf{a}^2$, which breaks the gauge symmetry down to $U(1)$. An interesting consequence of the inclusion of the θ -term in the 't Hooft-Polyakov action was pointed out by Witten in [22]. A necessary condition for gauge invariance of the partition function is

$$\exp \left\{ 2\pi i \left(\frac{Q}{e} - \frac{\theta e}{8\pi^2} \frac{M}{\hbar} \right) \right\} = 1 \quad \Rightarrow \quad \frac{Q}{e} - \frac{\theta e}{8\pi^2} \frac{M}{\hbar} = n_e \quad \text{with } n_e \in \mathbf{Z}. \quad (30)$$

Allowing M to be an integer multiple of the 't Hooft-Polyakov monopole charge

$$M = n_m \left(\frac{4\pi\hbar}{e} \right), \quad (31)$$

(which differs from (12) by a factor of 2 because the gauge group is $SO(3)$ and there are no spinor representations allowed) we see that

$$\frac{Q}{e} = n_e + n_m \frac{\theta}{2\pi} \quad (32)$$

with n_e and n_m integers, the electric and magnetic quantum numbers (for the 't Hooft-Polyakov monopole $n_e = 0$ and $n_m = -1$). This reduces to $Q = n_e e$ when $\theta = 0$ as before. If $\frac{\theta}{2\pi} = \frac{q}{p}$ is rational, with q and p mutually prime integers, then there exists the possibility of a magnetically charged particle with zero electric charge, when $(n_m, n_e) = (-p, q)$. Furthermore, since q and p are mutually prime, it is an elementary result of number theory that there also exist integers (n_m, n_e) such that

$$n_e p + n_m q = 1 \quad \text{and hence} \quad Q = \frac{e}{p}, \quad (33)$$

i.e. the Witten effect allows for fractionally charge particles. We shall see in §4 that the fractionally charged particles observed in the quantum Hall effect are a 2+1-dimensional analogue of this.

The masses of dyons in this model are related to the vacuum expectation value of the Higgs field and the electric and magnetic quantum numbers, in particular the masses \mathcal{M} are bounded below by the Bogomol'nyi bound [16, 32]

$$\mathcal{M}^2 \geq \frac{\mathbf{a}^2}{e^2} (Q^2 + M^2). \quad (34)$$

3 Duality in $\mathcal{N} = 2$, $SU(2)$ SUSY Yang-Mills

So far the discussion has been semi-classical: Dirac considered a quantised particle moving in a fixed classical background field and modular transformations are unlikely to be useful in the full theory of QED when coupled electromagnetic and matter fields are quantised. Nevertheless Seiberg and Witten showed in 1994 [4] that in fully quantised $\mathcal{N} = 2$ supersymmetric Yang-Mills theory, where the non-abelian gauge group is broken to $U(1)$, modular transformations of the complex coupling, or at least a subgroup thereof, are manifest in the vacuum structure. In essence supersymmetry is such a powerful constraint that, in the low-energy/long-wavelength limit of the theory, a remnant of the semi-classical modular action survives in the full quantum theory.

Consider $SU(2)$ Yang-Mills in 4-dimensional Minkowski space with global $\mathcal{N} = 2$ supersymmetry and, in the simplest case, no matter fields.³ The field content is then the $SU(2)$ gauge potential, A_μ ; two Weyl spinors (winos or gluinos) in a doublet of $\mathcal{N} = 2$ supersymmetry, both transforming under the adjoint representation of $SU(2)$, and a single complex scalar ϕ , again in the adjoint representation. The bosonic part of the action is

$$S = \int dx^4 \left\{ -\frac{1}{2g^2} \text{tr}(F_{\mu\nu} F^{\mu\nu}) + \frac{\theta}{32\pi^2} \varepsilon^{\mu\nu\rho\sigma} \text{tr}(F_{\mu\nu} F_{\rho\sigma}) + \frac{1}{g^2} \text{tr} \left((D_\mu \phi)^\dagger D^\mu \phi - \frac{1}{2} [\phi^\dagger, \phi]^2 \right) \right\}. \quad (35)$$

The fermionic terms (not exhibited explicitly here) are dictated by supersymmetry and involve Yukawa interactions with the scalar field but it is crucial that supersymmetry relates all the couplings and the only free parameters are the gauge coupling g and the topological susceptibility θ , all other couplings are in fact determined by g . Just as in electrodynamics these parameters can be combined into a single complex parameter which, in the conventions of [4], is

$$\tau = \frac{\theta}{2\pi} + i \frac{4\pi}{g^2}. \quad (36)$$

(Note that the imaginary part of τ is defined here to be twice that in §2 — this is because the 't Hooft-Polyakov monopole is central to the understanding of $N = 2$ SUSY Yang-Mills and it has twice the fundamental magnetic charge (12).) The definition (36) allows equations (31) and (32) to be combined into the single complex charge

$$Q + iM = g(n_e + n_m \tau) \quad (37)$$

and this is a central charge of the supersymmetry algebra. Semi-classical arguments would then imply modular symmetry (29) on τ [4], but we shall see that quantum effects break full modular symmetry, though a sub-group still survives in the full quantum theory.

Classically the Higgs potential is minimised by any constant ϕ in the Lie algebra of $SU(2)$ such that $[\phi^\dagger, \phi] = 0$, and since we can always rotate ϕ by a globally well-defined gauge transformation, we can take $\phi = \frac{1}{2} \mathbf{a} \sigma_3$ with $\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ the usual Pauli matrix and \mathbf{a} a complex constant with dimensions of mass. The classical vacua are thus highly degenerate and can be parameterised by \mathbf{a} , a non-zero \mathbf{a} breaks $SU(2)$ gauge symmetry down to $U(1)$ (we are still free to rotate around the σ_3 direction) and W^\pm bosons acquire a mass proportional to \mathbf{a} , leaving one $U(1)$ gauge boson (the photon) massless. At the same time the fermions and the scalar field also pick up masses proportional to \mathbf{a} except for the superpartners of the photon which are protected by supersymmetry and also remain massless. At the special point $\mathbf{a} = 0$ the gauge symmetry

³There is by now a number of good reviews of $\mathcal{N} = 2$ SUSY, see for example [33, 34].

is restored to the full $SU(2)$ symmetry of the original theory. Supersymmetry protects this vacuum degeneracy so that it is not lifted and is still there in the full quantum theory. At low energies, much less than the mass \mathbf{a} for a generic value of \mathbf{a} , the only relevant degrees of freedom in the classical theory are the massless $U(1)$ gauge boson and its superpartners (quantum effects modify this). Dyons with magnetic and electric quantum numbers (n_m, n_e) have a mass \mathcal{M} that, in $\mathcal{N} = 2$ SUSY, saturates the Bogomol'nyi bound (34) which in the conventions used here gives

$$\mathcal{M}^2 = \frac{2}{g^2} |\mathbf{a}(Q + iM)|^2 \quad (38)$$

(some factors of 2 differ from the original 't Hooft-Polyakov framework). Motivated by semi-classical modular symmetry this can be written, using (37), as

$$\mathcal{M}^2 = 2|\mathbf{a}(n_e + n_m\tau)|^2 = 2|n_e\mathbf{a} + n_m\mathbf{a}_D|^2 \quad (39)$$

where $\mathbf{a}_D := \tau\mathbf{a}$ so the 't Hooft-Polyakov monopole has mass $\mathcal{M}^2 = 2|\mathbf{a}_D|^2$. Under $\Gamma(1)$ τ transforms as

$$\gamma(\tau) = \frac{a\tau + b}{c\tau + d} = \frac{a\mathbf{a}_D + b\mathbf{a}}{c\mathbf{a}_D + d\mathbf{a}} \quad (40)$$

so

$$\begin{pmatrix} \mathbf{a}_D \\ \mathbf{a} \end{pmatrix} \rightarrow \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \mathbf{a}_D \\ \mathbf{a} \end{pmatrix} = \begin{pmatrix} a\mathbf{a}_D + b\mathbf{a} \\ c\mathbf{a}_D + d\mathbf{a} \end{pmatrix}. \quad (41)$$

The dyon mass \mathcal{M} is then invariant if at the same time (n_m, n_e) is transformed in the opposite way,

$$\begin{pmatrix} n_m \\ n_e \end{pmatrix} \rightarrow \begin{pmatrix} d & -c \\ -b & a \end{pmatrix} \begin{pmatrix} n_m \\ n_e \end{pmatrix} = \begin{pmatrix} dn_m - cn_e \\ -bn_m + an_e \end{pmatrix}. \quad (42)$$

In particular, under the analogue of Dirac's electric-magnetic duality: $\tau \rightarrow \tau_D = -1/\tau$ and $(\mathbf{a}_D, \mathbf{a}) \rightarrow (-\mathbf{a}, \mathbf{a}_D)$ and $(n_m, n_e) \rightarrow (-n_e, n_m)$. The central charge (37) is not invariant under these transformations

$$Q + iM \rightarrow \frac{1}{c\tau + d}(Q + iM) \quad (43)$$

unless $Q + iM$ is zero or infinite.

When the model is quantised there is a number of very important modifications to the classical picture:

- $\tau \rightarrow \tau_D = -1/\tau$ is not a symmetry of the quantum theory.
- The point $\mathbf{a} = 0$, where full $SU(2)$ symmetry is restored in the classical theory, is inaccessible in the quantum theory. The dual strong coupling point $\mathbf{a}_D = 0$ is accessible.
- The effective low-energy coupling runs as a function of \mathbf{a} , and since \mathbf{a} is related to a mass this is somewhat analogous to the Callan-Symanzik running of the QED coupling. The effective gauge coupling only runs for energies greater than $|\mathbf{a}|$, for energies less than this the running stops and at low energies it becomes frozen at its value $g^2(\mathbf{a})$. Using standard asymptotic freedom arguments in gauge theory, the low energy effective coupling $g^2(\mathbf{a})$ thus decreases at large \mathbf{a} and increases at small \mathbf{a} . In fact the topological susceptibility θ also runs with \mathbf{a} (as a consequence of instanton effects [4, 35]) and this can be incorporated into an \mathbf{a} -dependence of τ , $\tau(\mathbf{a})$. At large \mathbf{a} the imaginary part of τ , $\Im\tau$, becomes large while at small \mathbf{a} it is small.

- It is no longer true that $\tau = \mathbf{a}_D/\mathbf{a}$, in the quantum theory \mathbf{a} and \mathbf{a}_D are no longer linearly related but instead

$$\tau = \frac{\partial \mathbf{a}_D}{\partial \mathbf{a}}. \quad (44)$$

The classical mass formula (39) is only true in the quantum theory in the second form

$$\mathcal{M}^2 = \frac{2}{g^2} |n_e \mathbf{a} + n_m \mathbf{a}_D|^2. \quad (45)$$

A better, gauge invariant, parameterisation of the quantum vacua is given by $u = tr \langle \phi^2 \rangle$. For weak coupling (large \mathbf{a}) $u \approx \frac{1}{2} \mathbf{a}^2$, but $\langle \phi^2 \rangle \neq \langle \phi \rangle \langle \phi \rangle$ for strong coupling (small \mathbf{a}). By making a few plausible assumptions, including

- The low energy effective action is analytic except for isolated singularities (holomorphicity is intimately linked with supersymmetry);
- The number of singularities is the minimum possible compatible with stability of the theory ($\Im \tau > 0$),

Seiberg and Witten argued [4] that in the quantum theory the strong coupling regime $g^2 \approx 0$ is associated not with $\mathbf{a} = 0$ but instead with two points in the complex u -plane, $u = \pm \Lambda^2$ where Λ is, by definition, the mass scale at which the gauge coupling diverges. Furthermore they found an explicit expression for the full low energy effective action and argued that new massless modes appear at the singular points $u = \pm \Lambda^2$, in addition to the photon and its superpartners. These new massless modes are in fact dyons, with the magnetic charge coming from topologically non-trivial aspects of the classical theory (monopoles). Since $g \rightarrow \infty$ when $u = \pm \Lambda^2$, τ is real at these points. For example the point $\tau = 0$ is associated with $u = \Lambda^2$ and the dyons have zero electric charge, they are in fact simple 't Hooft-Polyakov monopoles with $n_m = -1$ ($M = -\frac{4\pi\hbar}{g}$) and $n_e = 0$, also $\mathbf{a}_D = 0$ at this point so $\mathcal{M} = 0$ from (45). The point $\tau = 1$ is associated with $u = -\Lambda^2$ and the dyons have $n_e = 1$ and $n_m = -1$, so $Q = 0$, and $\mathbf{a} = \mathbf{a}_D$ so again $\mathcal{M} = 0$.

The full modular group $\Gamma(1)$ is not manifest in the quantum theory, rather Seiberg and Witten showed that the relevant map is (29) with both b and c constrained to be even. This is a subgroup of the full modular group, denoted $\Gamma(2)$ in the mathematical literature, and it is generated by

$$\mathcal{T}^2 : \tau \rightarrow \tau + 2 \quad \text{and} \quad \mathcal{F}^2 : \tau \rightarrow \frac{\tau}{1 - 2\tau}, \quad (46)$$

where $\mathcal{F}^2 = \mathcal{S}^{-1} \mathcal{T}^2 \mathcal{S}$.⁴ u itself is invariant under $\Gamma(2)$, but $\tau(u)$ is a multi-branched function of u .

The explicit form of τ is given in [33] as

$$\tau = i \frac{K'}{K} + 2n \quad (47)$$

where n is an integer, $K = K(k)$ and $K' = K(k')$ are standard elliptic integrals with $k^2 = \frac{2}{1+u/\Lambda^2}$ and $k'^2 = 1 - k^2$ (see for example [36] for properties of elliptic integrals).

Seiberg and Witten's $\Gamma(2)$ action commutes with the scaling flow as u is varied. Taking the logarithmic derivative of $\tau(u)$ with respect to u , and imposing $ad - bc = 1$, we see that

$$u \frac{d\gamma(\tau)}{du} = \frac{1}{(c\tau + d)^2} u \frac{d\tau}{du}. \quad (48)$$

⁴We define $\mathcal{F} = \mathcal{S}^{-1} \mathcal{T} \mathcal{S} : \tau \rightarrow \frac{\tau}{1-\tau}$.

Meromorphic functions $\tau(u)$ satisfying (48) are well studied in the mathematical literature and are called modular forms of weight -2 .

For Seiberg and Witten's expression for $\tau(u)$ it was shown in [37, 38, 39, 40] that

$$-u \frac{d\tau}{du} = \frac{1}{2\pi i} \left(\frac{1}{\vartheta_3^4(\tau)} + \frac{1}{\vartheta_4^4(\tau)} \right), \quad (49)$$

where

$$\vartheta_3(\tau) = \sum_{n=-\infty}^{\infty} e^{i\pi n^2 \tau} \quad \text{and} \quad \vartheta_4(\tau) = \sum_{n=-\infty}^{\infty} (-1)^n e^{i\pi n^2 \tau} \quad (50)$$

are Jacobi ϑ -functions.⁵ Scaling functions for $\mathcal{N} = 2$ SUSY Yang-Mills are also constructed in [41]. At weak coupling, $g^2 \rightarrow 0$, $\tau \rightarrow i\infty$, $\vartheta_3(\tau) \rightarrow 1$ and $\vartheta_4(\tau) \rightarrow 1$ so

$$u \frac{d\tau}{du} \approx \frac{a}{2} \frac{d\tau}{da} \rightarrow \frac{i}{\pi} \quad (51)$$

which is the correct behaviour of the $\mathcal{N} = 2$ one-loop β -function for the gauge coupling,

$$a \frac{dg}{da} \approx -\frac{g^3}{4\pi^2}. \quad (52)$$

At weak coupling (large \mathbf{a}) \mathbf{a} is proportional to the gauge boson mass and this flow can be interpreted as giving the Callan-Symanzik β -function for the gauge bosons (or equivalently the winos because of supersymmetry) in the asymptotic regime. This interpretation is however not valid for finite \mathbf{a} for two reasons: firstly the statement that \mathbf{a} is proportional to the gauge boson mass is only valid at weak coupling and secondly because (49) diverges at strong coupling, $g \rightarrow \infty$ where $\tau \rightarrow 0$, [40]. The latter difficulty can be remedied by defining a different scaling function which is still a modular form of weight -2 . Seiberg and Witten's expression for $\tau(u)$ can be inverted to give $u(\tau)$ [38, 39, 40]

$$\frac{u}{\Lambda^2} = \frac{\vartheta_3^4 + \vartheta_4^4}{\vartheta_3^4 - \vartheta_4^4} \quad (53)$$

which is invariant under $\Gamma(2)$ modular transformations [39, 42]. Equation (49) can therefore be multiplied by any ratio of polynomials in u and we still have a modular form of weight -2 .⁶ The singularity in $u \frac{d\tau}{du}$ at $\tau = 0$ (where $u = \Lambda^2$) can be removed by multiplying by $(1 - u/\Lambda^2)^k$ with k some positive integer, without introducing any new singularities or zeros, which is the minimal assumption. Indeed the β -function at $\tau = 0$ can be calculated using one-loop perturbation theory in the dual coupling $\tau_D = -1/\tau$, [34], and it is shown in [44] that one obtains the correct form of the Callan-Symanzik β -function (up to a factor of 2), for winos at $\tau = i\infty$ and for monopoles at $\tau = 0$, using $k = 1$,

$$-(1 - u/\Lambda^2)u \frac{d\tau}{du} = \frac{1}{\pi i} \frac{1}{\vartheta_3^4(\tau)}. \quad (54)$$

However this is still not correct as there is a second singularity at $\tau = 1$, where $u = -\Lambda^2$. It is further argued in [44] that the correct form of the Callan-Symanzik β -function at $\tau = i\infty$, $\tau = 0$ and $\tau = 1$, up to a constant factor, is obtained by using the scaling function

$$-\left(1 - \frac{u}{\Lambda^2}\right) \left(1 + \frac{u}{\Lambda^2}\right) u \frac{d\tau}{du} = \frac{2}{\pi i} \frac{1}{(\vartheta_3^4(\tau) + \vartheta_4^4(\tau))} \quad (55)$$

⁵Definitions and relevant properties of the Jacobi functions can be found in the mathematics literature, for example [36].

⁶This point was stressed in [38]. See, for example, theorem 4.3.4 of [43].

and this is the minimal choice in the sense that it has the fewest possible singular points. The flow generated by (55) is shown in figure 1, there are fixed points on the real axis at $\tau = q/p$ where the massless dyons have quantum numbers $(n_m, n_e) = (-p, q)$. Odd p corresponds to attractive fixed points in the IR direction and even p to attractive fixed points in the UV direction ($p = 0$ corresponds to the original weakly coupled theory, $\tau = i\infty$). The repulsive singularities at $\tau = \frac{n+i}{2}$ with n an odd integer occur for $u = 0$ and are the quantum vestige of the classical situation were full $SU(2)$ symmetry would be restored.

One point to note is that, since the scaling function (55) is symmetric under $u \rightarrow -u$, which is equivalent to $\tau \rightarrow \tau + 1$, the full symmetry of the scaling flow is slightly larger than $\Gamma(2)$, it is generated by \mathcal{F}^2 and \mathcal{T} and corresponds to matrices γ such that c in (29) is even. This group is often denoted by $\Gamma_0(2)$.

Observe also that there are semi-circular trajectories linking some of the IR attractive fixed points with odd monopole charge. These can all be obtained from the semi-circular arch linking $\tau = 0$ and $\tau = 1$ by the action of some $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(2)$. Then

$$\tau_1 = q_1/p_1 = \gamma(0) = \frac{b}{d} \quad (56)$$

and

$$\tau_2 = q_2/p_2 = \gamma(1) = \frac{a+b}{c+d} \quad (57)$$

from which $b = q_1$, $d = p_1$ and $q_2 = a + b$, $p_2 = c + d$. Hence

$$q_2 p_1 - q_1 p_2 = 1, \quad (58)$$

since $ad - bc = 1$, giving a selection rule for transitions between IR attractive fixed points as the masses u are varied. This is clearly related to the Dirac-Schwinger-Zwanziger quantisation condition (16).

When matter in the fundamental representation is included the picture changes in detail, but is similar in structure [5]. In particular different subgroups of $\Gamma(1)$ appear. To anticipate the notation let $\Gamma_0(N) \subset \Gamma(1)$ denote the set of matrices with integral entries and unit determinant $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ such that $c = 0 \pmod{N}$ and let $\Gamma^0(N) \subset \Gamma(1)$ denote those with $b = 0 \pmod{N}$. These are both subgroups of $\Gamma(1)$: $\Gamma_0(N)$ being generated by \mathcal{T} and $\mathcal{S}^{-1}\mathcal{T}^N\mathcal{S}$ while $\Gamma^0(N)$ is generated by \mathcal{T}^N and $\mathcal{S}^{-1}\mathcal{T}\mathcal{S}$.

Now consider $\mathcal{N} = 2$ supersymmetric $SU(2)$ Yang-Mills theory in 4-dimensional Minkowski space with N_f flavours in the fundamental representation of $SU(2)$. The low energy effective action for $0 < N_f < 4$ was derived in [5] ($N_f = 4$ is a critical value for which the quantum theory has $\beta(\tau) = 0$ and the theory is conformally invariant). Because matter fields in the fundamental representation of $SU(2)$ can have half-integral electric charges it is convenient to re-scale τ by a factor of two and define

$$\tau' = \frac{\theta}{\pi} + \frac{8\pi i}{g^2}. \quad (59)$$

Thus

$$\gamma(\tau) = \tau + 1 \quad \Rightarrow \quad \gamma(\tau') = 2\gamma(\tau) = \tau' + 2 \quad (60)$$

and

$$\gamma(\tau) = \frac{\tau}{1-2\tau} \quad \Rightarrow \quad \gamma(\tau') = 2\gamma(\tau) = \frac{\tau'}{1-\tau'}, \quad (61)$$

so $\Gamma_0(2)$ acting on τ is equivalent to $\Gamma^0(2)$ acting on τ' .

The quantum modular symmetries of the scaling function acting on τ' are

$$\begin{aligned} N_f = 0, & \quad \Gamma^0(2) \\ N_f = 1, & \quad \Gamma(1) \\ N_f = 2, & \quad \Gamma_0(2) \\ N_f = 3, & \quad \Gamma_0(4) \end{aligned} \tag{62}$$

and explicit forms of the corresponding modular β -functions are given in [45]. For $N_f = 1$ and $N_f = 3$ the group is the same as the symmetry group acting on the effective action while for $N_f = 0$ and $N_f = 2$ it is larger, due to the \mathbf{Z}_2 action on the u -plane. Note the maximal case of $N_f = 1$ where the full modular group is manifest at the quantum level, in this case the duality transformation $\tau' \rightarrow -1/\tau'$ is a symmetry and $\tau' = i$ is a fixed point. The flow for $N_f = 0$ is shown in figure 1 and that for $N_f = 1$ is shown in figure 2, the other cases can be found in [45].

4 Duality and the Quantum Hall Effect

Modular symmetry manifests itself in the quantum Hall effect (QHE) in a manner remarkably similar to that of $\mathcal{N} = 2$ supersymmetric Yang-Mills. But while $\mathcal{N} = 2$ supersymmetry is not generally believed to have any direct relevance to the spectrum of elementary particles in Nature, the quantum Hall effect is rooted in experimental data and its supremely rich structure was not anticipated by theorists. The first suggestion that the modular group may be related to the QHE was in [2] but, as we shall see below, the wrong sub-group was identified in this earliest attempt.

The quantum Hall effect is a phenomenon associated with 2-dimensional semiconductors in strong transverse magnetic fields at low temperatures, so that the thermal energy is much less than the cyclotron energy $\hbar\omega_c$, with ω_c the cyclotron frequency. It requires pure samples with high charge carrier mobility μ so that the dimensionless product $B\mu$ is close to unity. For reviews see *e.g.* [46, 47, 48]. Basically passing a current I through a rectangular 2-dimensional slice of semi-conducting material requires maintaining a voltage parallel to the current (the longitudinal voltage V_L). The presence of a magnetic field B normal to the sample then generates a transverse voltage (the Hall voltage V_H). Two independent conductivities can therefore be defined: a longitudinal, or Ohmic, conductivity σ_L and a transverse, or Hall, conductivity σ_H , along with the associated resistivities, ρ_L and ρ_H . The classical Hall relation is

$$B = en\rho_H \quad \Rightarrow \quad \sigma_H B = -J_e^0 \tag{63}$$

(with $J_e^0 = en$ and \mathbf{n} the density of mobile charge carriers) and σ_H is inversely proportional to B at fixed \mathbf{n} . In the quantum Hall effect σ_H is quantised as $1/B$ is varied keeping \mathbf{n} and T fixed (or varying \mathbf{n} keeping B and T fixed) and increases in a series of sharp steps between very flat plateaux. At the plateaux σ_L vanishes and it is non-zero only for the transition region between two adjacent plateaux. In 2-dimensions conductivity has dimensions of e^2/h and, in the first experiments, [49] $\sigma_H = n\frac{e^2}{h}$ was an integral multiple of e^2/h at the plateaux (the integer QHE) though in later experiments [50] it was found that σ_H could also be a rational multiple of e^2/h , $\sigma_H = \frac{q}{p}\frac{e^2}{h}$ where p is almost always an odd integer (from now in this section on we shall adopt units in which $e^2/h = 1$). The different quantum Hall plateaux are interpreted as being different phases of a 2-dimensional electron gas and transitions between the phases can be induced by varying the external magnetic field, keeping the charge carrier density constant. An example of experimental data is shown in figure 3, taken from [51], where the Hall resistivity (ρ_{xy}) and the Ohmic resistivity (ρ_{xx}) for a sample are plotted as functions of the transverse magnetic field, in

units of h/e^2 . The Hall resistivity is monotonic, showing a series of steps or plateaux, while the Ohmic resistivity shows a series of oscillations with deep minima, essentially zero, when ρ_{xy} is at a plateau, and a series of peaks between the Hall plateaux.

Conductivity is actually a tensor

$$J_i = \sigma_{ij} E^j \quad (64)$$

with σ_{xx} and σ_{yy} the longitudinal conductivities in the x and y directions and $\sigma_{xy} = -\sigma_{yx} = \sigma_H$ the Hall conductivity associated with the magnetic field (for an elementary discussion of the Hall effect see [52]). From now on we shall assume an isotropic medium with $\sigma_{xx} = \sigma_{yy} = \sigma_L$. Using complex coordinates $z = x + iy$ the conductivity tensor for an isotropic 2-dimensional medium can be described by a single complex conductivity

$$\sigma := \sigma_H + i\sigma_L. \quad (65)$$

Note that Ohmic conductivities must be positive for stability reasons, so σ is restricted to the upper half-complex plane. The resistivity tensor is the inverse of the conductivity matrix, in complex notation $\rho_H + i\rho_L = \rho = -1/\sigma$.

In [15] it was argued that the following transformations

$$\mathcal{T} : \sigma \rightarrow \sigma + 1 \quad \text{and} \quad \mathcal{F}^2 : \sigma \rightarrow \frac{\sigma}{1 - 2\sigma} \quad (66)$$

map between different quantum Hall phases of a spin polarised sample. These transformations are known as the Law of Corresponding States for the quantum Hall effect.⁷

The \mathcal{T} transformation is interpreted as being due to shifting Landau levels by one, *e.g.* by varying the magnetic field keeping \mathbf{n} fixed. The philosophy here is that a full Landau level is essentially inert and does not affect the dynamics of quasi-particles in higher Landau levels. The number of filled Landau levels is the integer part of the filling factor $\nu = \frac{nh}{eB}$ and quasi-particles in a partially filled Landau level n , with n a non-negative integer, and filling factor $\nu = n + \delta\nu$, $0 \leq \delta\nu < 1$, will have similar dynamics to situations with the same $\delta\nu$ and any other n . In particular $n \rightarrow n + 1$ should be a symmetry of the quasi-particle dynamics. The situation here is similar to that of the periodic table of the elements were fully filled electron shells are essentially inert and do not affect the dynamics of electrons in higher shells — noble gases would then correspond to exactly filled Landau levels (the quantum Hall effect is in better shape than the periodic table of the elements however because all Landau levels have the same degeneracy whereas different electron shells have different degeneracies). Of course this assumes that there is no inter-level mixing by perturbations and this can never really be true for all n . To achieve large n for a fixed carrier density \mathbf{n} , for example, requires small B and eventually B will be so small that the inter-level gap, which is proportional to B , will no longer be large compared to thermal energies and/or Coulomb energies.

The \mathcal{F} transformation, known as flux-attachment, was used by Jain [53, 54], though only for Hall plateaux $\sigma_L = 0$, as a mapping between ground state wave-functions. Jain's mapping was associated with modular symmetry in [55] and with modular transformations of partition functions in [56]. It is related to the composite fermion picture of the QHE [47, 57, 58] where the effective mesoscopic degrees of freedom are fermionic particles bound to an even number of magnetic flux-tubes, the operation \mathcal{F}^2 attaches two units of flux to each composite fermion, maintaining their fermionic nature, in a manner that will be described in more detail below.

The response functions (*i.e.* the conductivities) in a low temperature 2-dimensional system can be obtained from a 2+1-dimensional field theory by integrating out all the microscopic

⁷Complex conductivities were not used in [15], but their results are more easily expressed in the notation used here.

physics associated with particles and/or holes and incorporating their contribution to the macroscopic physics into effective coupling constants. The classical Hall relation (63) can be derived from

$$\mathcal{L}_{eff}[A_0, J_e^0] = \sigma_H A_0 B + A_0 J_e^0, \quad (67)$$

where J_e^0 is the charge density of mobile charges only, not including the positive neutralising background of the ions. The co-variant version of this is

$$\mathcal{L}_{eff}[A, J_e] = \frac{\sigma_H}{2} \epsilon^{\mu\nu\rho} A_\mu \partial_\nu A_\rho + A_\mu J_e^\mu, \quad (68)$$

where $\mu, \nu, \rho = 0, 1, 2$ label 2+1 dimensional coördinates. In linear response theory the Hall conductivity here would be considered to be a response function, the response of the system to an externally applied electromagnetic field. Ohmic conductivity can be included by working in Fourier space (ω, \mathbf{p}) and introducing a frequency dependent electric permittivity. In a conductor the infinite wavelength electric permittivity has a pole at zero frequency, in a Laurent expansion

$$\epsilon(\omega, 0) = -i \frac{\sigma_L}{\omega} + \dots, \quad (69)$$

and all of the microscopic physics can be incorporated into response functions that modify the effective action for the electromagnetic field. In Fourier space, in the infinite wavelength, low-frequency limit, the modification is

$$\tilde{\mathcal{L}}_{eff}[A] \approx \frac{i\sigma_L}{4\omega} F^2 + \frac{\sigma_H}{4} \epsilon^{\mu\nu\rho} A_\mu F_{\nu\rho}, \quad (70)$$

where $F^2 = F_{\mu\nu}(-\omega) F^{\mu\nu}(\omega)$ etc. Of course the full dynamics is 3+1 dimensional, unlike the electrons the electric and magnetic fields are not confined to the 2-dimensional plane of the sample. What is being written here is the correction to the electromagnetic action due to the charges in the sample, from which response functions can be read off. The most relevant correction, in the renormalisation group sense, is the Chern-Simons term in (70). Naïvely assuming that there are no large anomalous dimensions arising from integrating out the microscopic degrees of freedom, the next most relevant term would be the usual Maxwell term. It is an assumption of the analysis in [15] that other terms, higher order in F and its derivatives, are not relevant.

Note that the right hand side of (70) is not real, an indication of the dissipative nature of Ohmic resistance, and non-local in time, again a feature of a conducting medium. Also we have used a relativistic notation and F^2 should really be split into $\mathbf{E}\cdot\mathbf{E}$ and $\mathbf{B}\cdot\mathbf{B}$ with different co-efficients (response functions). In the $\mathbf{p} = 0$, low-frequency limit of a conductor however, both response functions behave as $1/\omega$, the ratio is a constant and a relativistic notation can be used.⁸ Chern-Simons theories of the QHE have been considered by a number of authors [15, 59, 60, 61, 62, 63, 64, 65]. The presence of the F^2 term has been analysed from the general point of view of 3-dimensional conformal field theory in [66, 67, 68, 69].

For strong magnetic fields however (70) is not small and linear response theory cannot be trusted. This problem can be evaded by introducing what is called the “statistical gauge field”, a_μ , [58, 60, 70, 71], and then following the analysis of [15]. Consider a sample of material with N charge carriers with wave-function $\Psi(\mathbf{x}_1, \dots, \mathbf{x}_N)$. Perform a gauge transformation

$$\Psi(\mathbf{x}_1, \dots, \mathbf{x}_N) \rightarrow \Psi'(\mathbf{x}_1, \dots, \mathbf{x}_N) = e^{i\vartheta(\sum_{i<j} \phi_{ij})} \Psi(\mathbf{x}_1, \dots, \mathbf{x}_N) \quad (71)$$

where ϕ_{ij} is the angle related to the positions of particle i and particle j as shown in figure 4, in terms of complex coördinates, $\mathbf{z} = x + iy$, $e^{i\phi_{ij}} = \frac{\mathbf{z}_i - \mathbf{z}_j}{|\mathbf{z}_i - \mathbf{z}_j|}$. ϑ is a constant parameter and

⁸For finite ω and/or non-zero \mathbf{p} this would not be the case as the response functions for \mathbf{E} and \mathbf{B} would be different functions in general.

can change the statistics of the particles: for example if the particles are fermions, so that Ψ is anti-symmetric under interchange of any two particles i and j , *i.e.* when $\phi_{ij} \rightarrow \phi_{ij} + \pi$, then Ψ' is again anti-symmetric if $\vartheta = 2k$ is an even integer, Ψ' becomes symmetric if $\vartheta = 2k + 1$ is an odd integer.

For co-variant derivatives, $(-i\hbar\nabla - eA)_\mu$ acting on Ψ becomes $(-i\hbar\nabla - eA - a)_\mu$ acting on Ψ' under (71) where

$$a_\alpha(\mathbf{x}_i) = \hbar\vartheta \sum_{j \neq i} \nabla_\alpha^{(i)} \phi_{ij} = -\hbar\vartheta \sum_{j \neq i} \epsilon_{\alpha\beta} \frac{(\mathbf{x}_i - \mathbf{x}_j)^\beta}{|\mathbf{x}_i - \mathbf{x}_j|^2}, \quad (\alpha, \beta = 1, 2). \quad (72)$$

As long as the particles have a repulsive core, so $\mathbf{x}_i \neq \mathbf{x}_j$ for $i \neq j$ and the potential a is a pure gauge, but if the particles can co-incide there is a vortex singularity in a and

$$e^{\beta\alpha} \nabla_\beta^{(i)} a_\alpha(\mathbf{x}_i) = \hbar\vartheta \sum_j \delta^{(2)}(\mathbf{x}_i - \mathbf{x}_j) \rightarrow \hbar\vartheta \mathbf{n}(\mathbf{x}_i), \quad (73)$$

the last expression being the continuum limit of the discrete particle distribution. The statistical gauge field a_μ was introduced in the context of ‘‘anyons’’ in [60, 70]. Its relation with the charge carrier density (73) was pointed out in [71], where similarities between the statistics parameter ϑ and the vacuum angle QCD were noted and it was observed the binding of vortices to particles is a 2+1-dimensional version of ’t Hooft’s notion of oblique confinement in QCD [72], where a condensate of composite objects occurs.

The constraint (73) can be encoded into the dynamics, by introducing a Lagrange multiplier a_0 , and then made co-variant by modifying (68) to read

$$\mathcal{L}_{eff}[A, a, J^e] = \frac{s}{2} \epsilon^{\mu\nu\rho} A_\mu \partial_\nu A_\rho + \left(A + \frac{a}{e}\right)_\mu J_e^\mu - \frac{1}{2\vartheta e^2} \epsilon^{\mu\nu\rho} a_\mu \partial_\nu a_\rho, \quad (74)$$

(using units with $\frac{e^2}{\hbar} = 1$) where J_e^μ is the current generated by the matter fields Ψ' . Note that the co-efficient of the A_μ Chern-Simons term has been changed from σ_H to a new parameter s , this is because quantum effects modify the Hall co-efficient and identifying σ_H with s is premature at this stage. Following [15] is now argued that integrating out Ψ' in (74) to get an effective action for A and a can only produce terms that depend on the combination $A' := A + a/e$ and gauge invariance restricts the allowed terms.⁹ Define the field strength for A' as usual

$$F'_{\mu\nu} = \partial_\mu A'_\nu - \partial_\nu A'_\mu \quad (75)$$

and that of a as

$$f_{\mu\nu} = \partial_\mu a_\nu - \partial_\nu a_\mu. \quad (76)$$

Then the effective action, in Fourier space in the long-wavelength $\mathbf{p} = 0$ limit, will be of the form

$$\tilde{\mathcal{L}}_{eff}[A, a] = \frac{s}{4} \epsilon^{\mu\nu\rho} A_\mu F_{\nu\rho} - \frac{\Pi_L(\omega)}{4} F'^2 + \frac{\Pi_H(\omega)}{4} \epsilon^{\mu\nu\rho} A'_\mu F'_{\nu\rho} - \frac{1}{4\vartheta e^2} \epsilon^{\mu\nu\rho} a_\mu f_{\nu\rho}, \quad (77)$$

where the response functions Π_L and Π_H cannot be calculated but are *assumed* to give conductivities in the $\omega \rightarrow 0$ limit, *i.e.* it is assumed that Π_L has a pole while Π_H is finite at $\omega = 0$,

$$i\omega \Pi_L(\omega) \xrightarrow{\omega \rightarrow 0} \sigma_L, \quad \Pi_H(\omega) \xrightarrow{\omega \rightarrow 0} \sigma_H. \quad (78)$$

⁹Note that Euclidean signature was used in [15] while we use Minkowski signature here.

Integrating out Ψ' will in general produce many more terms in the effective action than shown in (77), higher order powers in F' and its derivatives, with unknown co-efficients, but a further assumption is that these are less relevant, in the renormalisation group sense, than the Π_L and Π_H terms exhibited here. This does require a leap of faith however — while it is certain that there are no anomalous dimensions associated with the Chern-Simons term, $\epsilon A'F'$ since it is topological, it is by no means obvious that integration of Ψ' will not produce large anomalous dimensions associated with operators like $(F'^2)^2$, for example, that might render them more relevant than F'^2 . It is an assumption in [15] that this does not happen. Note however that it *has* been assumed that there is a large anomalous dimension associate with F'^2 — this operator is not naïvely marginal in 2+1-dimensions, but the assumption of a pole in Π_L renders it marginal. The effective action (77) is actually conformal [66]. One further modification of the effective action used in [15] is to allow for a change in the effective charge of the matter fields Ψ' , $e^* = \eta e$. This is easily incorporated into (77) by redefining A' to be $A' = \eta A + a/e$ everywhere (the units are not affected, $h = e^2$).

The \mathcal{F}^2 and \mathcal{T} transformations in (66) can now be derived by integrating a out of the effective action (77), which is quadratic in a and so the integration is Gaussian. This leads to a new effective action for A alone, with different conductivities,

$$\tilde{\mathcal{L}}'_{eff}[A] = \frac{i\sigma'_L}{4\omega} F^2 + \frac{\sigma'_H}{4} \epsilon^{\mu\nu\rho} A_\mu F_{\nu\rho} \quad (79)$$

where σ'_L and σ'_H are best expressed in terms of the complex conductivity $\sigma' = \sigma'_H + i\sigma'_L$ as a fractional linear transformation

$$\sigma' = \frac{a\sigma + b}{c\sigma + d} \quad (80)$$

and one finds [15]

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \frac{1}{\eta} \begin{pmatrix} \eta^2 + s\vartheta & s \\ \vartheta & 1 \end{pmatrix} := \gamma(\eta, s, \vartheta). \quad (81)$$

The analysis of was taken further in [77] and extended beyond the regime of linear response in [78].

Note that $\det \gamma = 1$ so the most general transformation, for arbitrary s , ϑ and η , is an element of $Sl(2; \mathbf{R})$. However a general element is not a symmetry, for example if ϑ is not an integer then the two conductivities describe charge carriers which are anyons with different statistics. Also if ϑ is an odd integer we have mapped between bosonic and fermionic charge carriers and this is not a symmetry, though this was the map described in the analysis in [15] where σ was associated with a bosonic system, which could form a Bose condensation, and σ' was associated with a quantum Hall system with fermionic charge carriers (there is no suggestion here that a supersymmetric theory necessarily underlies the quantum Hall effect).

Quantum mechanics requires that the parameters η , s and ϑ in (81) must necessarily be quantised for the map to be a symmetry. For example $\eta = 1$, when the charge carriers are pseudo-particles with $e^* = e$, and $\vartheta = 0$ gives

$$\gamma(1, s, 0) = \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix} \quad (82)$$

and we only expect this to be a symmetry when s is an integer (Landau level addition). In particular $\gamma(1, 1, 0)$ gives the \mathcal{T} transformation of (66). The flux attachment transformation, \mathcal{F}^2 in (66), follows from the fact that $\vartheta = 2k$ should be a symmetry for any integral k , with $\eta = 1$ and $s = 0$. In particular

$$\gamma(1, 0, -2) = \begin{pmatrix} 1 & 0 \\ -2 & 1 \end{pmatrix} \quad (83)$$

gives \mathcal{F}^2 .

We can now map between different Hall plateaux by combining the operations of adding statistical flux and Landau level addition. For example if we start from a plateau with $\sigma = 1$ and perform a (singular) gauge transformation with $\vartheta = 2$, which is \mathcal{F}^{-2} , then we transform to a new phase with $\sigma = 1/3$.

Note that a general transformation requires $\eta \neq 1$, although it is always rational. For example the Jain series of fractional quantum Hall states [73] can be generated by starting from $\sigma = 0$ and mapping to Landau level q using \mathcal{T}^q , where q is a positive integer. Now add $2k$ units of statistical flux using \mathcal{F}^{-2k} , the resulting transformation is

$$\mathcal{F}^{-2k}\mathcal{T}^q = \begin{pmatrix} 1 & 0 \\ 2k & 1 \end{pmatrix} \begin{pmatrix} 1 & q \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & q \\ 2k & 2kq + 1 \end{pmatrix} = \gamma \left(\frac{1}{2kq + 1}, \frac{q}{2kq + 1}, \frac{2k}{2kq + 1} \right). \quad (84)$$

The new conductivity has

$$\sigma' = \frac{q}{p} \quad \text{with} \quad p = 2kq + 1 \quad \text{and} \quad \eta = \frac{1}{p}, \quad (85)$$

so the pseudo-particles carry not only $2k$ vortices of statistical flux but also electric charge

$$e^* = \frac{e}{p} \quad (86)$$

with p odd, *i.e.* fractional charge. This is precisely analogous to the Witten effect described in §2. Fractional charges of $e/3$ for $q = k = 1$ were predicted in [74] and have been confirmed by experiment [75, 76]. The analysis of [15] was generalised to the regime of non-linear response in [78].

The model of the Hall plateaux espoused in [15] is that one has charge carriers which are fermions (electrons or holes) interacting strongly with the external field. By attaching an odd number $2k + 1$ of statistical flux units to each fermion the resulting composite particles are bosons. By choosing k appropriately it can be arranged that the effect of the external magnetic field is almost cancelled by the statistical gauge field and the composite bosons behave almost as free particles. Being bosons they can condense to form a superconducting phase, with a mass gap, and this explains the stability of the quantum Hall plateaux for the original fermions. This map between bosons and fermions is of course not a symmetry, nevertheless it is a useful way of looking at the physics. An alternative view is Jain's composite fermion picture [47, 57, 58], which is a symmetry. Both the composite fermion and the composite boson picture are useful, but in either case fermions lie at the heart of the physics, as indicated by the antisymmetry of the Laughlin ground state wave-functions.

The transformations (66) acting on the complex conductivity map between different quantum Hall phases, clearly they generate the group $\Gamma_0(2)$. As mentioned above the relevance of the modular group to the QHE was anticipated by Wilczek and Shapere [2], though these authors focused on a different subgroup of $\Gamma(1)$, one generated by

$$\mathcal{S} : \sigma \rightarrow -1/\sigma \quad \text{and} \quad \mathcal{T}^2 : \sigma \rightarrow \sigma + 2, \quad (87)$$

and denoted by Γ_θ here and by $\Gamma_{1,2}$ in [2], and the experimental data on the QHE do not bear this out. For example Γ_θ has a fixed point at $\sigma = i$ there is no such fixed point in the experimental data for the quantum Hall effect. We shall return to Γ_θ below in the context of 2-dimensional superconductors. Lütken and Ross [12] observed, even before [15], that the quantum Hall phase diagram in the complex conductivity plane bore a striking resemblance to the structure of moduli space in string theory, at least for toroidal geometry, and postulated that $\Gamma(1)$ was relevant to the quantum Hall effect. In [13, 14] the sub-group $\Gamma_0(2)$ was identified

as being one that preserves the parity of the denominator and therefore likely to be associated with the robustness of odd denominators in the experimental data.

Among the assumptions that go into the derivation of (66) from (70) are that the temperature is sufficiently low and that the sample is sufficiently pure, but unfortunately the analyses in [12, 13, 15, 78] are unable to quantify exactly what “sufficiently” means. Basically one must simply assume that (70) contains the most relevant terms for obtaining the long wavelength, low frequency response functions, but obviously this is not always true even at the lowest temperatures, for example it is believed that, at zero temperature, a Wigner crystal will form for filling fractions below about 1/7 and (70), being rotationally invariant, cannot allow for this.¹⁰

Of course $\Gamma_0(2)$ is not a symmetry of all of the physics, after all the conductivities differ on different plateaux, nevertheless it is a symmetry of some physical properties. The derivation of the $\Gamma_0(2)$ action in [15] required performing a Gaussian integral about a fixed background, different backgrounds give different initial conductivities, but in each case the dynamics contributing to the fluctuations are the same and the dynamics of the final system have the same form as that of the initial one. This motivates the suggestion [80, 81, 82] that the scaling flow, which is governed by the fluctuations, should commute with $\Gamma_0(2)$, *i.e.* although $\Gamma_0(2)$ is not a symmetry of all the physics it is a symmetry of the scaling flow. Physically the scaling flow of the QHE can be viewed as arising from changing the electron coherence length l , *e.g.* by varying the temperature T with $l(T)$ a monotonic function of T [83, 84]. Define a scaling function by

$$\Sigma(\sigma, \bar{\sigma}) := l \frac{d\sigma}{dl}. \quad (88)$$

In general one expects σ to depend on various parameters, such as the temperature T , the external field B , the charge carrier density \mathbf{n} and the impurity density \mathbf{n}_I . If \mathbf{n} and \mathbf{n}_I are fixed then $\sigma(B, T)$ becomes a function of B and T only. The scaling hypothesis of [84] suggests that, at low temperatures, σ becomes a function of a single scaling variable.

Now for any $\gamma \in \Gamma(1)$, $\gamma(\sigma) = \frac{a\sigma+b}{c\sigma+d}$ with $ad - bc = 1$, so

$$\Sigma(\gamma(\sigma), \gamma(\bar{\sigma})) = \frac{1}{(c\sigma + d)^2} \Sigma(\sigma, \bar{\sigma}), \quad (89)$$

this is always true, by construction, and does not represent a symmetry.

Demanding that $\Gamma_0(2)$ is a symmetry of the QHE flow we immediately get very strong predictions concerning quantum Hall transitions. Firstly one can show that any fixed point of $\Gamma_0(2)$, with $\sigma_{xx} > 0$, must be fixed point of the scaling flow (though not *vice versa*), [80]. To see this assume that the scaling flow commutes with the action of $\Gamma_0(2)$ and let σ_* be a fixed point of $\Gamma_0(2)$, *i.e.* there exists a $\gamma \in \Gamma_0(2)$ such that $\gamma(\sigma_*) = \sigma_*$. If σ_* were not a scaling fixed point, we could move to an infinitesimally close point $\delta f(\sigma_*) \neq \sigma_*$ with an infinitesimal scaling transformation, δf . Assuming $\gamma(\delta f(\sigma_*)) = \delta f(\gamma(\sigma_*)) = \delta f(\sigma_*)$ then implies that $\delta f(\sigma_*)$ is also left invariant by γ . But, for $\Im\sigma > 0$, the fixed points of $\Gamma_0(2)$ are isolated and there is no other fixed point of $\Gamma_0(2)$ infinitesimally close to σ_* . Hence $\delta f(\sigma_*) = \sigma_*$ and σ_* must be a scaling fixed point. Fixed points σ_* of $\Gamma_0(2)$ with $\Im\sigma > 0$ are easily found by solving the equation $\gamma(\sigma_*) = \sigma_*$: there is none with $\Im\sigma > 1/2$: there is one at $\sigma_* = \frac{1+i}{2}$, indeed a series with $\sigma_* = \frac{n+i}{2}$ for all integral n , and of course the images of these under $\Gamma_0(2)$, leading to a fractal structure as one approaches $\Im\sigma = 0$. Many of these fixed points have been seen in experiments [80].

Experiments show that fixed points between plateaux correspond to second order phase transitions between two quantum Hall phases and modular symmetry implies that scaling exponents should be the same at each fixed point, a phenomenon known as *super-universality* [85, 86]. The idea here is that the transition between two quantum Hall plateaux is a quantum phase

¹⁰Fractions as low as 1/9 have been reported at finite temperature [79].

transition [87] and, at low temperatures, σ becomes a function of the single scaling variable $\frac{\Delta B}{T^\kappa}$ with a scaling exponent κ , where $\Delta B = B - B_c$ with B_c the magnetic field at the critical point. While there is strong evidence for super-universality with $\kappa \approx 0.44$ [85], [88], there are some experiments which seem to violate it [89]. This may be due to the suggestion that there exists a marginal operator at the critical point [90], but an alternative explanation, that the scaling function might be a modular form, is proposed in [91].

A second striking consequence of $\Gamma_0(2)$ symmetry of the scaling flow is a selection rule for quantum Hall transitions which can be derived in exactly the same way as in the discussion around (58) in §3. Given that the integer transition $\sigma : 2 \rightarrow 1$ is observed we conclude that $\gamma(2) \rightarrow \gamma(1)$ should also be possible. Let $\gamma(1) = \frac{q_1}{p_1}$ and $\gamma(2) = \frac{q_2}{p_2}$ where q_1 and p_1 are mutually prime, as are q_2 and p_2 . Then, with $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$,

$$\frac{q_1}{p_1} = \frac{a+b}{c+d} \quad \text{and} \quad \frac{q_2}{p_2} = \frac{2a+b}{2c+d} \quad (90)$$

so

$$q_1 = a + b, \quad p_1 = c + d, \quad q_2 = 2a + b \quad \text{and} \quad p_2 = 2c + d.$$

Since $ad - bc = 1$ we obtain the selection rule [80]

$$q_2 p_1 - q_1 p_2 = 1. \quad (91)$$

This selection rule is well borne out by the experimental data in figure 3. In spin-split samples any two adjacent well-formed plateaux, with no unresolved sub-structure between them, obey this rule.¹¹

If we further assume that the *only* fixed points of the scaling flow are the fixed points of $\Gamma_0(2)$ then, with a few extra reasonable assumptions, the topology of the flow diagram is completely determined and it is exactly the same as in figure 1 for $\mathcal{N} = 2$ SUSY. The whole upper-half plane can be covered by starting with the infinite strip of width 1 above the semi-circular arch spanning 0 and 1 in figure 1 (called the fundamental domain of $\Gamma_0(2)$ and shown in figure 5, [43]) and acting on it with elements of $\Gamma_0(2)$. IR fixed points have $\tau = q/p$ corresponding to fermionic charge carriers which can be viewed, by turning [15] around, as bosons with p units of statistical flux attached, where p is odd.

Guided by experiment, the reasonable assumptions are [91]:

- Rational numbers q/p with odd p are attractive, since they are experimentally so stable;
- The flow comes down vertically from the points at $\sigma = \sigma_H + i\infty$ *i.e.* that σ_H does not flow at high temperatures (weak coupling [91, 92])

then the topology forces the fixed points with $\Im\sigma > 0$ to be saddle points and the only flow possible is compatible with the topology of figure 1. The precise form of the flow in figure 1 can be deformed, but the fixed points cannot be moved. An even stronger statement can be made with one further assumption. When a system is symmetric under particle-hole interchange one has symmetry under change of the sign of σ_H , which is $\sigma \rightarrow 1 - \bar{\sigma}$ for the complex conductivity. This puts a reality condition on $\Sigma(\sigma, \bar{\sigma})$ which, combined with the mathematical properties of invariant functions of $\Gamma_0(2)$, can be used to show that the boundary of the fundamental domain in figure 5, and its images under $\Gamma_0(2)$ must always be flow lines, [93]. Any deformation of figure 1 must

¹¹Note the point labelled 11/9 in figure 3 is not compatible with the adjacent plateau at 9/7 according to this selection rule. However 11/9 is merely a kink in the flow which has probably been identified wrongly. The selection rule implies that this kink is most likely be the precursor of the 14/11 plateau and that there is unresolved structure between 14/11 and the integer plateau at 1.

therefore leave invariant not only the fixed points but also the vertical lines above integers on the real axis and the semi-circles spanning rational numbers on the real line with odd denominators, (as well as the images of these under $\Gamma_0(2)$) — only the specific shape of the flow lines inside the fundamental domain, and their images under $\Gamma_0(2)$, can be deformed. In particular $\Gamma_0(2)$ symmetry provides a remarkably robust derivation of the well-established experimental semi-circle law. In many experiments the transition between two plateaux does follow a semi-circle in the upper-half complex conductivity plane to a very high degree of accuracy, figure 6 for example is taken from [94]. (Note however that the critical point, as indicated by B_c in figure 6, is not at $\sigma = (1+i)/2$ as would be predicted by $\Gamma_0(2)$ symmetry. This will be discussed below.) The scaling hypothesis of [84, 85] described above suggests that, at low temperatures, σ becomes a function of the scaling variable $\frac{\Delta B}{T^\kappa}$ and the flow is then forced onto one of these semi-circles as $T \rightarrow 0$. At a microscopic level the semi-circle law has been derived in one specific model [95, 96], but modular symmetry provides a very robust derivation, [93], valid for any model which satisfies the above assumptions, and is therefore much more general than any specific model. Any deviation from a semi-circular transition is an experimental signal that the sample under investigation is not symmetric under particle-hole interchange. The general topology of the flow in figure 1, at least for the integer QHE, was predicted by Khmel'nitskii [83], though the normalisation of the vertical axis was not determined in that analysis and there was no extension to the fractional case.¹²

The specific flow for figure 1 was determined by supplementing the above assumptions with an additional constraint, that the scaling function $\Sigma(\sigma)$ should be a meromorphic function in the argument σ .¹³ This makes $\Sigma(\sigma)$ a modular form of weight -2 and the analytic form that approaches the stable fixed points on the real axis most rapidly is then exactly the same as that of $\mathcal{N} = 2$ SUSY Yang-Mills in the previous section, up to an undetermined constant, namely

$$\Sigma(\sigma) = \frac{2}{\pi i} \frac{1}{(\vartheta_3^4(\sigma) + \vartheta_4^4(\sigma))}, \quad (92)$$

and this gives the flow plotted in figure 1. To date there is no physical reason for assuming that Σ should be independent of $\bar{\sigma}$, it is only motivated by mathematical analogy with SUSY Yang-Mills, but it does have the advantage of giving an explicit form for the scaling function that can be visualised. Any deformation away from this by including a non-meromorphic component is constrained by the considerations above and cannot change the topology. In fact the flow obtained using the meromorphic ansatz does give remarkably good agreement with experiment and the comparison is plotted in figures 7 and 8, taken from [97, 98] (in figure 7 the Landau levels are spin-degenerate and this has the effect of doubling the conductivity — see below and footnote 14).

In summary, modular symmetry $\Gamma_0(2)$ applied to the quantum Hall effect for spin-split samples leads to the following predictions:

- Universal critical points are predicted at $\sigma_* = \frac{1}{2}(1+i)$ and its images under $\Gamma_0(2)$. Critical exponents must be the same for all fixed points which are related by $\Gamma_0(2)$.
- Exact flow lines in the σ plane can be derived from $\Gamma_0(2)$ invariance plus invariance under particle-hole symmetry: $\sigma \rightarrow 1 - \bar{\sigma}$. In particular the semi-circle law is a consequence of $\Gamma_0(2)$ plus particle-hole symmetry.
- Flow in the infrared is towards the real axis, terminating on the real axis at attractive fixed points at odd-denominator fractions. Even-denominator fractions form repulsive fixed points of the flow.

¹²A similar topology appears in the flow of one-dimensional clock models [8], but there is no obvious connection with the modular group in these models.

¹³Strictly speaking it is meromorphic in the variable $q = e^{i\pi\sigma}$ rather than in σ itself.

- The selection rule $|p_1q_2 - p_2q_1| = 1$ for allowed transition between $\sigma = q_1/p_1$ and $\sigma = q_2/p_2$.

For QHE samples that are not fully spin polarised one expects that the Landau levels can come in adjacent pairs and \mathcal{T} in (66) should be replaced by \mathcal{T}^2 , giving the generators of $\Gamma(2)$ rather than $\Gamma_0(2)$, and this modifies the experimental consequences. $\Gamma(2)$ actually has no fixed points above the real axis, so one cannot make any predictions about the position of the second order phase transition between two plateaux, but if there is particle-hole symmetry one still has the semi-circle law and the fixed point must lie somewhere on the semi-circle, [99]. An experimental analysis of the relation between modular symmetry and Zeeman splitting is given in [100, 101, 102] and the results support the interpretation in [99]. Figure 9 is reproduced from [102] and represents a sample where the spin splitting is small, significantly less than the Landau level splitting but still non-negligible in the low Landau levels where the magnetic field is large, and $\Gamma_0(2)$ symmetry is broken to $\Gamma(2)$, so the critical point, Q in the figure, is not at the top of the semi-circle. Nevertheless particle-hole symmetry is obeyed reasonably well, as manifest by the close approximation to a semi-circle. In the same sample, at lower magnetic fields and higher Landau levels the spin-splitting becomes negligible and the sample becomes spin-degenerate. In this situation $\Gamma_0(2)$ symmetry again applies, but with the conductivity doubled because of the spin degeneracy in each Landau level. The critical point in the $\nu = 2 \rightarrow \nu = 4$ transition is indeed at the top of the semi-circle in the conductivity plane, as reported in [102].¹⁴ It is possible that the explanation of the fact that B_c is not at the top of the semi-circle in figure 6 is due to the spins being poorly split, but there is another possible interpretation for this $\nu : 1 \rightarrow 0$ transition [99]. This particular transition is a vertical line in the complex ρ -plane above the point $\rho = 1$ and a re-scaling of ρ_{xx} can move the critical point, which should be at $\rho_{xx} = 1$ if $\Gamma_0(2)$ symmetry holds, to any value of ρ_{xx} and hence to any point on the corresponding semi-circle in the σ -plane. If the determination of the scale of ρ_{xx} in [94] has an error then figure 6 is still compatible with $\Gamma_0(2)$ symmetry. For the particular case of the $\nu : 1 \rightarrow 0$ transition any re-scaling of ρ_{xx} still gives a semi-circle in the σ -plane, but this is not true for other transitions. The observation that the critical point is not at $\sigma = (3+i)/2$ for the $\nu : 2 \rightarrow 1$ transition in [102], as shown in figure 9, cannot be explained by a re-scaling of the Ohmic resistivity and therefore is interpreted as a breaking of $\Gamma_0(2)$ symmetry down to $\Gamma(2)$ because the Landau levels, while not exactly degenerate, are not well split.

The group $\Gamma(2)$ was also analysed, from the point of view of its action on ground state wave-functions rather than on complex conductivities, in [103] and on complex conductivities in [104, 105].

The group $\Gamma_0(2)$ is the subgroup of the full modular group that is relevant for systems with fermionic charge carriers in a strong perpendicular magnetic field with the spins well split. Building on the work of [15], and following a suggestion in [106], it was shown in [77] that $\Gamma_0(2)$ should be the group relevant to 2-dimensional systems involving spin polarised fermionic quasi-particles while Wilczek and Shapere's group Γ_θ should be relevant to 2-dimensional systems involving bosonic quasi-particles. If the effective charge carriers are bosonic, *e.g.* 2-dimensional superconductors, one can obtain predictions from the fermionic case by using the flux attachment transformation to add an odd number of vortices to the quasi-particles. For the case of a single unit of flux this is equivalent to conjugating $\Gamma_0(2)$ by $\mathcal{F} = \mathcal{S}^{-1}\mathcal{T}\mathcal{S}$. Since $\mathcal{F}^{-1}\mathcal{F}^2\mathcal{F} = \mathcal{F}^2$ and $\mathcal{F}^{-1}\mathcal{T}\mathcal{F} = \mathcal{F}^{-2}\mathcal{S}$ the resulting group is generated by \mathcal{S} and \mathcal{F}^2 , or equivalently \mathcal{S} and \mathcal{T}^2 , and is the group Γ_θ , [2, 78, 77]. Note that the \mathcal{S} transformation in (87) requires $\vartheta = -s = \pm\eta$ with $\eta \rightarrow \infty$ in (81). The fixed point $\sigma = i$ of \mathcal{S} is associated with a phase transition from an insulator to a superconductor [107] and a superconductor would have a double pole in its response function, associated with spontaneous symmetry breaking and requiring a mass term

¹⁴Because of the spin degeneracy the transformations are $\sigma \rightarrow \sigma + 2$ and $\sigma \rightarrow \frac{\sigma}{1-\sigma}$, *i.e.* $\Gamma_0(2)$ acting on $\sigma/2$ is equivalent to $\Gamma^0(2)$ acting on σ , just as for $N_f = 3$ in SUSY.

$m^2 A^2$ in the effective action. A direct derivation of \mathcal{S} transformation, in conjunction with particle-hole symmetry, for a 2-dimensional superconductor was given in [108]. Γ_θ consists of elements γ of the form (28) with either a, d both odd and b, c both even, or *vice versa*. In this case the predictions are different:

- Universal critical points are predicted for the flow at the fixed points for transitions between stable phases, $\sigma_* = i$ and its images under Γ_θ .¹⁵
- The critical exponents at all fixed points related by Γ_θ must all be the same.
- Exact flow lines in the σ plane are immediate consequences of Γ_θ invariance and particle-hole symmetry, [77]. The results are again semi-circles or vertical lines in the σ plane, implying a semi-circle law for these bosonic systems.
- For nonzero magnetic fields the flow as the temperature is reduced is towards the real axis, terminating on the attractive fixed points $\sigma = p/q$ with pq even (as opposed to having q odd, as was the case for fermions). Fractions with odd pq are repulsive fixed points.
- There is a selection rule that allowed fractions p_2/q_2 can be obtained from p_1/q_1 only if $|p_1 q_2 - p_2 q_1| = 1$.

The resulting flow diagram for bosonic systems was given in [77] and a more detailed version is shown in figure 10. It has a fixed point at $\sigma_* = i$, as predicted by Fisher [107]. To date no 2-dimensional superconductors have been manufactured with a high enough mobility μ that μB is close to unity for sustainable magnetic fields, but it is predicted in [77] that a hierarchy with the above properties will be observed if such samples are ever manufactured.

5 Conclusions

Modular symmetry is a generalisation of the Dirac quantisation condition for charge in QED. Its mathematical foundation is strongest in supersymmetric systems, in particular supersymmetric Yang-Mills, but a more realistic system, with a wealth of experimental data to compare with, is the quantum Hall effect.

Effective degrees of freedom are composite objects carrying topologically non-trivial gauge field configurations (monopoles in SUSY Yang-Mills and vortices in the QHE).

For $\mathcal{N} = 2$ SUSY $SU(2)$ Yang-Mills the sub-groups of the modular group that are relevant for the scaling modular forms with different numbers of families are

$$\begin{aligned}
 N_f = 0, & & \Gamma^0(2) \\
 N_f = 1, & & \Gamma(1) \\
 N_f = 2, & & \Gamma_0(2) \\
 N_f = 3, & & \Gamma_0(4).
 \end{aligned} \tag{93}$$

For the quantum Hall effect the relevant sub-groups are

$$\begin{aligned}
 \text{spin-split samples,} & & \Gamma_0(2) \\
 \text{spin-degenerate samples,} & & \Gamma^0(2)
 \end{aligned} \tag{94}$$

$$\text{intermediate case,} \tag{95} \quad \Gamma(2).$$

¹⁵This statement is for bosonic charge carriers with the same electric charge as an electron — it becomes $\sigma_* = i\tilde{q}^2$ and its images if the bosonic charge carriers have charge \tilde{q} . In particular, the case $\tilde{q} = \pm 2$ applies if the bosons are Cooper pairs, such as considered in ref. [107].

Modular symmetry has important experimental consequences for the quantum Hall effect. The Dirac-Schwinger-Zwanziger quantisation condition manifests itself as a selection rule for transitions between allowed quantum Hall plateaux,

$$|qp' - q'p| = 1 \tag{96}$$

with Hall conductivities q/p and q'/p' in spin-split samples ($|qp' - q'p| = 2$ for spin-degenerate samples). Universal critical points in transitions between plateaux for spin-degenerate and spin-split samples are an unavoidable consequence of modular symmetry. Together with particle-hole symmetry, when it is present, modular symmetry immediately implies the semi-circle law for quantum Hall transitions — any deviation from the semi-circle law is most likely due to breaking of particle-hole symmetry in the sample. Fractional charges in the quantum Hall effect are related to the Witten effect in 3+1-dimensional $\mathcal{N} = 2$ supersymmetric Yang-Mills, where massive excitations can carry electric charge $1/p$ with p being the monopole number.

For bosonic pseudo-particles the group Γ_θ is predicted to be the relevant group and this gives a suite of predictions similar to the quantum Hall effect but differing in detail, as described in §4.

The connection between $\mathcal{N} = 2$ SUSY and the quantum Hall effect that is exposed by modular symmetry is not yet understood, at the moment it is merely at the level of an empirical observation that modular symmetry is relevant to scaling in both these systems. Constructing an explicit map relating one to the other remains an open problem to date. There is a known relation between $\mathcal{N} = 4$ SUSY and the quantum Hall effect, [109, 110, 111], which may prove to be useful starting point, perhaps via SUSY breaking, but since the perturbative β -function for $\mathcal{N} = 4$ SUSY vanishes there is no clear connection with scaling at present. There are however certain features which both $\mathcal{N} = 2$ SUSY and the QHE have in common. They all have only two relevant couplings: one, y , associated with the dynamical kinetic term, which must be positive for stability reasons; and one, x , associated with a topological term. These are combined into the complex parameter, $\tau = x + iy$, on the upper half-complex plane on which the modular group, or a sub-group thereof, acts.

An important feature that these systems have in common is the existence of topologically non-trivial field configurations, monopoles in SUSY Yang-Mills and vortices in the QHE, which can bind to the charge carriers to form pseudo-particles carrying magnetic charge: dyons in SUSY Yang-Mills and in the QHE fermions which can be viewed as bosons with an odd number of flux units attached. The rational nature of the attractive fixed points on the real axis q/p is then related to the magnetic charge p . The weak coupling regime, where perturbation theory might be expected to be useful, can be mapped by modular symmetry to the strong coupling regime, where much interesting physics lies.

It seems that modular symmetry is not just an accident of one type of system, or even a family of systems such as supersymmetric field theories, but in fact is a more general phenomenon and it may yet prove even more powerful in understanding the physics of strongly interacting systems.

- [1] B.P Dolan, *Mod. Phys. Lett.* **A21** (2006) 1567
- [2] A. Shapere and F. Wilczek, *Nuc. Phys.* **B320** (1989) 669
- [3] C. Montonen and D. Olive, *Phys. Lett.* **B72** (1977) 117
- [4] N. Seiberg and E. Witten, *Nuc. Phys.* **B426** (1994) 19; Erratum *ibid.* **B430** (1994) 169, [[hep-th/9407087](#)];
- [5] N. Seiberg and E. Witten, *Nuc. Phys.* **B431** (1994) 484, [[hep-th/9408099](#)]
- [6] J.L. Cardy and E. Rabinovici, *Nuc. Phys.* **B205** (1982) 1
- [7] J.L. Cardy, *Nuc. Phys.* **B205** (1982) 17
- [8] M. Asorey, J.G. Esteve and J. Salas, *Phys. Rev.* **B48** (1993) 3626
- [9] C.G. Callan and D. Freed, *Nucl. Phys.* **B374** (1992) 543, [[hep-th/9110046](#)]
- [10] C.G. Callan, A. Felce and D. Freed, *Nucl. Phys.* **B392** (1993) 551, [[hep-th/9202085](#)]
- [11] Wellington da Cruz, *Chaos, Solitons, Fractals* **17** (2003) 975, [[math-ph/029028](#)]; *Int. J. Mod. Phys.* **A18** (2003) 2213, [[cond-mat/0212567](#)]
- [12] C.A. Lütken and G.G. Ross, *Phys. Rev.* **B45** (1992) 11837;
- [13] C.A. Lütken and G.G. Ross, *Phys. Rev.* **B48** (1993) 2500
- [14] C.A. Lütken, *J. Phys.* **A26** (1993) L811; *Nuc. Phys.* **B396** (1993) 670
- [15] S. Kivelson, D-H. Lee and S-C. Zhang, *Phys. Rev.* **B46** (1992) 2223
- [16] J.M. Figueroa-O'Farrill, *Electromagnetic Duality for Children*, <http://www.maths.ed.ac.uk/~jmf/EDC.html>
- [17] P.A.M. Dirac, *Proc. Roy. Soc.* **A133** (1931) 60
- [18] T.T. Wu and C.N. Yang, *Phys. Rev.* **D12** (1975) 3845
- [19] K.A. Milton, *Rept. Prog. Phys.* **69** (2006) 1637, [[hep-ex/0602040](#)]
- [20] J. Schwinger, *Phys. Rev.* **144** (1966) 1987
- [21] D. Zwanziger, *Phys. Rev.* **D3** (1971) 880
- [22] E. Witten, *Phys. Lett.* **86B** (1979) 283
- [23] E. Witten, *Selecta. Math.* (NS) **1** (1995) 383, [[hep-th/9505186](#)];
- [24] E. Verlinde, *Nucl. Phys.* **B455** (1995) 211
- [25] A.M. Polyakov, *Phys. Lett.* **59B** (1975) 82
- [26] A.A. Belavin, A.M. Polyakov, A. Schwartz and Y. Tyupkin, *Phys. Lett.* **59B** (1975) 85
- [27] G. 't Hooft, *Phys. Rev. Lett.* **37** (1976) 8
- [28] C. Callan, R. Dashen and D. Gross, *Phys. Rev.* **63B** (1976) 334
- [29] R. Jackiw and C. Rebbi, *Phys. Rev. Lett.* **37** (1976) 334
- [30] G. 't Hooft, *Nuc. Phys.* **B79** (1974) 276
- [31] A.M. Polyakov, *JETP Lett.* **20** (1974) 194
- [32] E.B. Bogomol'nyi, *Soviet J. Nuc. Phys.* **24** (1976) 449
- [33] L. Alvarez-Gaumé and S.F. Hassan, *Fortsch. Phys.* **45** (1997) 159, [[hep-th/9701069](#)]
- [34] M.E. Peskin, *Duality in Supersymmetric Yang-Mills Theory*, [[hep-th/9702094](#)]
- [35] N. Dorey, V.V. Khoze and M.P. Mattis, *Phys. Rev.* **D54** (1996) 2921, [[hep-th/9603136](#)]
- [36] E.T. Whittaker and G.N. Watson, *A Course of Modern Analysis*, C.U.P. (1940)
- [37] J.A. Minahan and D. Nemeschansky, *Nucl. Phys.* **B468** (1996) 72, [[hep-th/9601059](#)]
- [38] A. Ritz, *Phys. Lett.* **B434** (1998) 54, [[hep-th/9710112](#)]

-
- [39] B.P. Dolan, *Phys. Lett.* **B418** (1998) 107, [hep-th/9710161]
- [40] G. Carlino, K. Konishi, N. Maggiore and N. Magnoli, *Phys. Lett.* **B455** (1999) 171, [hep-th/9902162]; K. Konishi, *Int. J. Mod. Phys.* **A16** (2001) 1861, [hep-th/0012122]
- [41] E. D'Hoker, I.M. Krichever and D.H. Phong, *Nuc. Phys.* **B489** (1997) 179, [hep-th/9609041]; *Nuc. Phys.* **B494** (1997) 89, [hep-th/9610156]
- [42] M. Matone, *Phys. Lett.* **B357** (1995) 342, [hep-th/9506102]
- [43] R.A. Rankin, *Modular Forms and Functions*, C.U.P. (1977)
- [44] B.P. Dolan, *N = 2 Supersymmetric Yang-Mills and the Quantum Hall Effect*, To appear in *Int. J. Mod. Phys. A*, [hep-th/0505138]
- [45] B.P. Dolan, *Nucl. Phys.* **B737** (2006) 153, [hep-th/0506088]
- [46] *The Quantum Hall Effect*, eds. R.E. Prange and S.M. Girvin, Springer (New York), (1987)
- [47] *Composite Fermions*, Ed. O. Heinonen, World Scientific (1998)
- [48] M. Stone, *Quantum Hall Effect*, World Scientific (1992)
- [49] K. von Klitzing, G. Dorda and M. Pepper, *Phys. Rev. Lett.* **45** (1980) 494
- [50] D.C Tsui, H.L. Stormer and A.C. Gossard, *Phys. Rev. Lett.* **48** (1982) 1559
- [51] D.C. Tsui, *Physica B* **164** (1990) 59
- [52] L. Landau and E. Lifschitz, *A Course in Theoretical Physics*, Vol. 8 (2nd Edition), Pergamon Press (1984)
- [53] J.K Jain, S.A. Kivelson and N. Trivedi, *Phys. Rev. Lett.* **64** (1990) 1927
- [54] J.K. Jain and V.J. Goldman, *Phys. Rev.* **B45** (1992) 1255
- [55] I. Kogan and R. Szabo, *Phys. Rev.* **B58** (1998) 7893, [cond-mat/9712233]
- [56] A. Cappelli and R. Zamba, *Nuc. Phys.* **B490** (1997) 595, [hep-th/9605127]
- [57] J.K. Jain, *Phys. Rev. Lett.* **63** (1989) 199; *Phys. Rev.* **B41** (1990) 7653
- [58] A. Lopez and E. Fradkin, *Phys. Rev.* **B44** (1991) 5246
- [59] S.M. Girvin, in [46]
- [60] D.P. Arovas, J.R. Schrieffer, F. Wilczek and A. Zee, *Nuc. Phys.* **B251** (1985) 117
- [61] S.C. Zhang, T.H. Hansson and S. Kivelson, *Phys. Rev. Lett.* **62** (1989) 82
- [62] J. Fröhlich and A. Zee, *Nuc. Phys.* **B364** (1991) 517
- [63] A.P. Balachandran and A.M. Srivastava, *Chern-Simons Dynamics and the Quantum Hall Effect*, [hep-th/9111006]
- [64] A. Balatsky and E. Fradkin, *Phys. Rev.* **B43** (1991) 10622
- [65] E. Fradkin, *Nuc. Phys.* **B474** [FS] (1996) 543
- [66] E. Witten, *SL(2, Z) Action On Three-Dimensional Conformal Field Theories With Abelian Symmetry*, [hep-th/0307041]
- [67] R.G. Leigh and A.C. Petkou, *JHEP* **0312** (2003) 020, [hep-th/0309177]
- [68] R. Zucchini, *Adv. Theor. Math. Phys.* **8** (2005) 895, [hep-th/0311143]
- [69] Ho-Ung Yee, *Phys. Lett.* **B598** (2004) 139, [hep-th/0402115]
- [70] F. Wilczek, *Phys. Rev. Lett.* **49** (1982) 957
- [71] S.M. Girvin and A.H. MacDonald, *Phys. Rev. Lett.* **58** (1987) 1252
- [72] G. 't Hooft, *Nucl. Phys.* **B190** [FS] (1981) 455
- [73] See for example the article by Jain and Kamilla in [47]

- [74] R.B. Laughlin, *Phys. Rev. Lett.*, **50** (1983) 1395
- [75] R. de-Picciotto *et al* , *Nature* **389** (1997) 162
- [76] L. Saminadayar *et al*, *Phys. Rev. Lett.* **79** (1997) 2526
- [77] C. Burgess and B.P. Dolan, *Phys. Rev.* **B63** (2001) 155309, [hep-th/0010246]
- [78] C. Burgess and B.P. Dolan, *Phys. Rev.* **B65** (2002) 155323, [cond-mat/0105621]
- [79] W. Pan *et al*, *Phys. Rev. Lett.* **88** (2002) 17680
- [80] B.P. Dolan, *J. Phys.* **A32** (1999) L243, [cond-mat/9805171]
- [81] N. Taniguchi, *Nonperturbative Renormalization Group Function for Quantum Hall Plateau Transitions Imposed by Global Symmetries* [cond-mat/9810334]
- [82] C. Burgess and A. Lütken, *Nuc. Phys.* **B500** (1997) 367, [cond-mat/9611118]; *Phys. Lett.* **B451** (1999) 365, [cond-mat/9812396]
- [83] D.E. Khmel'nitskii, *Pis'ma Zh. Eksp. Teor. Fiz* **38** (1983) 454, (*JETP Lett.* **38** (1983) 552)
- [84] A.M.M. Pruisken, *Phys. Rev. Lett.* **61** (1988) 1297
- [85] H.P. Wei, D.C. Tsui, M.A. Paalanen and A.M.M. Pruisken, *Phys. Rev. Lett.* **61** (1988) 1294
- [86] L. Engel, H.P. Wei, D.C. Tsui and M. Shayegan, *Surf. Sci.* **229** (1990) 13
- [87] S.L. Sondhi, S.M. Girvin, J.P. Carini and D. Shankar, *Rev. Mod. Phys.* **69** (1997) 315
- [88] D. Shahar, D. C. Tsui, M. Shayegan, E. Shimshoni and S. L. Sondhi, *Phys. Rev. Lett.* **79** (1997) 479, [cond-mat/9611011]
- [89] D. Shahar, M. Hilke, C.C. Li, D.C. Tsui, S.L. Sondhi and M. Razeghi, *Solid State Commun.* **107** (1998) 19, [cond-mat/9706045]
- [90] M.R. Zirnbauer, *Conformal Field Theory of the integer quantum Hall effect*, [hep-th/9905054]
- [91] B.P. Dolan, *Nucl. Phys.* **B554** (1999) 487, [cond-mat/9809294]
- [92] A.M.M. Pruisken, in [46]
- [93] C. Burgess, B.P. Dolan and Rim Dib, *Phys. Rev.* **B62** (2000) 15359, [cond-mat/9911476]
- [94] M. Hilke *et al.*, *Euro. Phys. Lett.* **46** (1999) 775, [cond-mat/9810217]
- [95] A.M. Dykhne and I.M. Ruzin, *Phys. Rev.* **B50** (1994) 2369
- [96] I. Ruzin and S. Feng, *Phys. Rev. Lett.* **74** (1995) 154
- [97] S.S. Murzin, M. Weiss, A.G.M. Jansen and K. Eberl, *Phys. Rev.* **B66** (2002) 233314, [cond-mat/0204206]
- [98] S.S. Murzin, S.I. Dorozhkin, D.K. Maude and A.G.M. Jansen, *Phys. Rev.* **B72** (2005) 195317, [cond-mat/0504235]
- [99] B.P. Dolan, *Phys. Rev.* **B62** (2000) 10278, [cond-mat/0002228]
- [100] C.F. Huang, Y.H. Chang, H.H. Cheng, C.-T. Liang and G.J. Hwang, *Physica* **E22** (2004) 232, [cond-mat/0404246];
- [101] D.R. Hang *et al*, *Microelectron. J.* **36** (2005) 469, [cond-mat/0508577]
- [102] C.F. Huang *et al*, *An experimental study on $\Gamma(2)$ modular symmetry on the quantum Hall system with a small spin-splitting*, [cond-mat/0609310]
- [103] M.A.I. Flohr, *On a New Universal Class of Phase Transitions and Quantum Hall Effect*, [hep-th/9412053]
- [104] Y. Georgelin, T. Masson and J.-C. Wallet, *J. Phys.* **A33** (1990) 39; *J. Phys.* **A30** (1997) 5065

-
- [105] Y. Georgelin and J-C. Wallet, *Phys. Lett.* **A224** (1997) 303;
 - [106] L.P. Pryadko, *Phys. Rev.* **B56** (1997) 6810
 - [107] M.P.A. Fisher, *Phys. Rev. Lett.* **65** (1990) 923
 - [108] S.-J. Rey and A. Zee, *Nucl. Phys.* **B352** (1991) 897
 - [109] D. Berenstein, *Phys. Rev.* **D71** (2005) 085001, [[hep-th/0409115](#)]
 - [110] H. Lin, O. Lunin and J. Maldacena *JHEP* **0410** (2004) 025, [[hep-th/0409174](#)]
 - [111] A. Ghodsi, A.E. Mosaffa, O. Saremi and M.M. Sheikh-Jabbari *Nucl. Phys.* **B729** (2005) 467, [[hep-th/0505129](#)]




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7





Figure 8



C. F. Huang et al.

Figure 9




Figure 10