

A Resummable β -Function for Massless QED

Oliver J. Rosten*

Dublin Institute for Advanced Studies, 10 Burlington Road, Dublin 4, Ireland

Within the set of schemes defined by generalized, manifestly gauge invariant exact renormalization groups for QED, it is argued that the β -function in the four dimensional massless theory cannot possess any nonperturbative power corrections. Consequently, the perturbative expression for the β -function must be resumable. This argument cannot be extended to flows of the other couplings or to the anomalous dimension of the fermions and so perturbation theory does not define a unique trajectory in the critical surface of the Gaussian fixed point. Thus, resumability of the β -function is not inconsistent with the expectation that a non-trivial fixed point does not exist.

PACS numbers: 11.10.Gh, 11.10.Hi

The resumability¹, or otherwise, of the perturbative series for the couplings and anomalous dimension(s) in some quantum field theory (QFT) is intimately related to the nonperturbative question of renormalizability. This is beautifully explained in [1] (see also [2]), and we here recall the main points. The formalism best suited to understanding such issues is the Exact Renormalization Group (ERG), which is essentially the continuous version of Wilson's RG [3, 4]. A fundamental ingredient of this approach is the implementation of a momentum cutoff, such that all modes above the cutoff scale are regularized. For the following discussion, we consider two cutoff scales. First, there is the bare scale, Λ_0 , which provides an overall cutoff to the theory. As we shall see, for nonperturbatively renormalizable theories, this scale is an artificial construction, and it is misleading to identify the action at this scale as a boundary condition that can be chosen, arbitrarily. (The same is not true for nonrenormalizable theories.) We now integrate out degrees of freedom between the bare scale and a lower, 'effective' scale, Λ . As we perform this procedure, the bare action evolves into the Wilsonian effective action, S_Λ , in such a way that the partition function stays the same. The Wilsonian effective action can be thought of as parametrizing the interactions relevant to the effective scale. The ERG equation states how the Wilsonian effective action changes with the effective scale.

One of the most important uses of the ERG equation is to find QFTs which are nonperturbatively renormalizable, in other words theories for which Λ_0 can be sent to infinity (this is called taking the continuum limit). Scale independent renormalizable theories follow immediately from fixed points of the ERG equation. To see this, suppose that we rescale all dimensionful quantities to dimensionless ones, by dividing by Λ raised to the appropriate scaling dimension. Now, fixed points of the ERG can be immediately identified with renormalizable theories: as a consequence of our rescalings, independence of Λ im-

plies independence of all scales; independence of all scales trivially implies independence of Λ_0 , and so obviously we can send Λ_0 to infinity!

Scale dependent renormalizable theories can be constructed by considering flows out of any of the fixed points, along the associated relevant / marginally relevant directions. The Wilsonian effective actions lying on these 'Renormalized Trajectories' (RTs) [3] are self-similar, meaning that all dependence on Λ appears only through the renormalized relevant / marginally relevant couplings and anomalous dimension(s).² Self-similarity implies renormalizability, since there is no explicit dependence on Λ/Λ_0 . Note that, along an RT, the theory is completely specified by the choice of fixed point, and the integration constants or 'rates' associated with the relevant / marginally relevant directions. In the limit $\Lambda \rightarrow \infty$, the theory sinks back into the appropriate fixed point. Thus, if we wish to consider the action at some arbitrarily high 'bare' scale, we must *compute* it using the flow equation, given our aforementioned choices. Indeed, the 'bare action' in this context is the perfect action [5] in the vicinity of the ultraviolet (UV) fixed point. This is in contrast to a nonrenormalizable trajectory, where we can simply choose some bare action, and use it as the boundary condition for the flow.

One of the benefits of viewing renormalization in this way is that, along RTs, we can compute directly in terms of renormalized quantities, without any mention of the bare scale or the bare action. To do this, we employ renormalization conditions for the relevant and marginally relevant couplings and the anomalous dimension(s) directly at the effective scale, Λ . So, if a non-trivial RT were to exist in QED (we are not claiming that one does in four dimensions, where the gauge coupling is marginally *irrelevant*, but one does in three dimensions) then we would define the gauge coupling—which we denote by g and not e to avoid later confusion— simply by

*Electronic address: orosten@stp.dias.ie

¹ Throughout this paper we have in mind Borel resumability, though our conclusions should not depend on this choice.

² Any masses are included in our definition of couplings.

writing the gauge kinetic term as

$$\frac{1}{g^2(\Lambda)} \int d^D x F_{\mu\nu} F_{\mu\nu},$$

at all scales. Note that we have scaled the coupling out of the gauge field. In the manifestly gauge invariant approach that we will later adopt, this will have the pleasant effect of guaranteeing that the gauge field does not suffer from field strength renormalization [6]. Throughout this paper we work in Euclidean space, so there is no distinction between upper and lower indices.

Let us now consider a massless theory, about which it is supposed that all we know is that its ERG trajectory lies in the critical surface of some fixed point. Since this trajectory is flowing *into* a fixed point, and we have not specified whether or not the trajectory happens to have emanated from some other fixed point in the UV, we do not know, a priori, whether the theory is renormalizable or not. To be concrete, we will suppose that this infrared fixed point is the Gaussian one, that this fixed point possesses a single marginally irrelevant coupling, g , and that there is a single anomalous dimension (just as there is in our manifestly gauge invariant approach to QED in four dimensions).

Let us now do perturbation theory in the vicinity of the Gaussian fixed point, within the critical surface. For reasons that will become apparent, we will attempt to write the action in self-similar form. Consequently, our renormalization conditions involve conditions for only the coupling, g , and the anomalous dimension, γ , specified at the scale Λ . We have assumed, temporarily, that no reference to the bare scale / bare action is necessary. Computing the full perturbative solution to the theory, we find that everything can be written in renormalized terms [6], defining an apparently unique, self-similar trajectory in the critical surface of the Gaussian fixed point. Were it really the case that this trajectory were both self-similar and unique, then this would suggest the existence of a UV fixed point, out of which an RT can be constructed that flows into the Gaussian fixed point. However, as emphasised in [1], this picture is generally false. In the specific case of scalar field theory in four dimensions, the perturbative series for the n -point couplings of the theory are not resummable, and so do not unambiguously define functions. The reason for this is UV renormalons³ (see [7] for a review of renormalons): perturbation theory by itself is not well defined, but must be supplemented by exponentially small terms which, in QED, take the form

$$\frac{\Lambda}{\Lambda_0} \sim e^{-1/2\beta_1 g^2(\Lambda)} + \dots, \quad (1)$$

³ Throughout this paper, we will loosely refer to a renormalon as *any* singularity of the Borel transform, rather than using the strict definition [7], which defines renormalons as those singularities related to large or small loop momentum behaviour.

where β_1 is the one-loop β -function and the ellipsis denotes higher order corrections. The left-hand side of this expression makes it immediately clear that self-similarity of our trajectory is violated: there is explicit dependence on Λ_0 . We can always write such power corrections in terms of g , as we have done on the right-hand side, but the prefactor will, of course, depend on Λ_0 . Within perturbation theory, we can blithely take the limit $\Lambda_0 \rightarrow \infty$, but at some point have to face up to the fact that this procedure is not well defined, if we hope to draw reliable nonperturbative conclusions.

In light of this discussion, the main result of this paper is rather unexpected: it will be demonstrated, in massless QED in four dimensions that, given a particular definition of the coupling, the perturbative series for the β -function *cannot be supplemented by terms of type (1) and its generalizations*. Since our ERG equation is perfectly well defined, and since we can choose a perfectly well defined boundary condition (bare action) we deduce that the perturbative β -function must be resumable. Nevertheless, our earlier conclusions are still intact, since our argument does not apply to the other couplings of the theory or to the anomalous dimension of the fermions. Consequently, perturbation theory still does not specify a unique, self-similar trajectory within the critical surface of the Gaussian fixed point, and so there is no suggestion that a UV fixed point exists.

Note, though, that matters could be much more interesting in the Wess-Zumino model. First, we note that all couplings belonging to the superpotential are protected from flowing by the nonrenormalization theorem. Secondly, it seems as though we can apply the arguments of this paper to show that the perturbative series for the anomalous dimension is resumable. (By scaling the field strength renormalization out of the two-point vertex, we can induce a flow of the superpotential and so relate the β -function of the three-point coupling to the anomalous dimension. It looks like the arguments applying to the β -function in this paper go through similarly in the Wess-Zumino model.) Finally, following [8], we can demonstrate that the flow of the dressed, exact n -point vertices can be written in terms of the apparently resumable anomalous dimension [for an example of a dressed two-point function, see (6), below]. Moreover, the relationship between the dressed vertices and the Wilsonian effective action vertices can be straightforwardly inverted [8]. This suggests the existence of a (well defined) self-similar trajectory in the critical surface, which would indicate the presence of a UV fixed point. Work on this is underway [9].

That terms of the type (1) are precluded comes about as follows. The key is to express the β -function as a ratio of two other functions [see (10), below]. Now, there is no reason to suppose that each of these two functions cannot, separately, possess contributions of the form (1). However, for reasons that we will precisely spell out later, any such contributions must exactly cancel each other.

The definition of our QED coupling is defined through

our choice of ERG. In this paper, we use the framework of generalized ERGs [10, 11], which can be used to furnish a manifestly gauge invariant formulation of QED [6] and even non-Abelian gauge theories [11–15]. The essence of this approach is as follows. As stated already, a necessary ingredient of the ERG equation is that the partition function is invariant under the flow. Consequently, given some set of fields, φ , we can define the family of ERGs to which Polchinski's equation [16] belongs according to [10]

$$-\Lambda \partial_\Lambda e^{-S[\varphi]} = \int_x \frac{\delta}{\delta\varphi(x)} \left(\Psi_x[\varphi] e^{-S[\varphi]} \right), \quad (2)$$

where the Λ derivative is performed at constant φ , any Lorentz indices etc. have been suppressed and we have written S_Λ as just S . It is the total derivative on the right-hand side ensures that the partition function $Z = \int \mathcal{D}\varphi e^{-S}$ is invariant under the flow.

The functional, Ψ , parametrizes a general Kadanoff blocking [17] in the continuum, for which we take the following form [11]:

$$\Psi_x = \frac{1}{2} \{ \hat{\Delta}^{\varphi\varphi}(x, y) \} \frac{\delta\Sigma}{\delta\varphi(y)}, \quad (3)$$

where it is understood that we sum over all the elements of the set of fields φ . Whilst we will leave the blocking procedure largely unspecified, there are certain general requirements that must be satisfied. Crucially, blocking must take place only over a localized patch, ensuring that each infinitesimal RG step is free of IR divergences.

We now describe each of the components on the right-hand side of (3). First, there are the ERG kernels, $\hat{\Delta}^{\varphi\varphi}$, which are generally different for each of the elements of φ . Each kernel incorporates a cutoff function which provides UV regularization. The notation $\{\hat{\Delta}\}$ denotes a covariantization of the kernel which may be necessary, depending on the symmetries of the theory. Indeed, it is apparent from (2) and (3) that the kernel essentially ties together two functional derivatives at points x and y ; in gauge theory, we can covariantize this statement by using e.g. straight Wilson lines between these two points. In practice, we leave any necessary covariantization unspecified, demanding only that it satisfies general requirements [11–13]. The remaining ingredient in (3) is $\Sigma \equiv S - 2\hat{S}$, where \hat{S} is the seed action [6, 11, 13, 18, 19]. Whereas we solve the flow equation for the Wilsonian effective action, the seed action serves as an input and, given our choice (3) and a choice of cutoff function(s), parametrizes the remaining freedom in how modes are integrated out along the flow.

The constraint that Ψ corresponds to a local blocking transformation translates into the requirement that the seed action leads to convergent momentum integrals and that the seed action and (covariantized) cutoff functions have all orders derivative expansions. In turn, this guarantees that the Wilsonian effective action vertices have a derivative expansion, also, this being a property that we will exploit, later.

The final point to make about (2) is a subtle one: it might be necessary to include some unphysical fields in the set φ , in order to properly implement a UV cut-off. Indeed, this is precisely the case in the manifestly gauge invariant ERG formulation of QED that we employ, where covariantization of the cutoff functions is not sufficient to completely regularize the theory: it is necessary to include Pauli-Villars (PV) partners for the fermions. (This is due to the well known result that covariant higher derivatives fail to regularize a set of one loop divergences.) Consequently, the field content for our manifestly gauge invariant ERG for QED comprises the gauge field, A_μ , a fermion field, ψ , and an unphysical commuting spinor field, χ , which is given a mass at the effective scale (it is obviously trivial to generalize to extra flavours). To be completely clear: when we loosely refer to QED, we strictly mean regularized theories of an Abelian vector field, coupled to fermions, whose effective action in the vicinity of the Gaussian fixed point is that of QED, to excellent approximation.

The precise details of the set-up can be found in [6], but we will not need them here. Rather, for our purposes, we need only consider the flow equation for the various *vertex coefficient functions* (i.e. all fields and symmetry factors having been stripped off), which has a generic diagrammatic form, largely independent of the details of the set-up and even the precise field theory being considered [6, 14, 19].

Given the aforementioned field content, we substitute (3) into (2), perform the Λ -derivative on the left-hand side and identify terms with the same numbers of fields [6, 11–14, 20]. Before doing this, we scale the coupling out of the covariant derivative, for reasons mentioned earlier. The rescaling causes $S \rightarrow S/g^2$ and, in contrast to some previous works [11, 13, 14, 21], we choose to similarly redefine the seed action. Thus, defining $\Sigma_g \equiv g^2(S - 2\hat{S})$, the diagrammatic flow equation for the vertex coefficient functions is shown in figure 1 [6].

$$\left(-\Lambda \frac{d}{d\Lambda} + \sum_{\phi \in \{f\}} \gamma^{(\phi)} \right) \left[\textcircled{S} \right]^{\{f\}} = \frac{1}{2} \left[\begin{array}{c} \textcircled{\Sigma_g} \\ \bullet \\ \textcircled{S} \end{array} - \textcircled{\Sigma_g} \right]^{\{f\}}$$

FIG. 1: The diagrammatic form of the QED flow equation for vertices of the Wilsonian effective action.

The first term on the left-hand side represents the flow of all independent Wilsonian effective action vertex coefficient functions corresponding to the set of fields, $\{f\}$. Since the Λ -derivative strikes just a vertex coefficient function—all fields having been stripped off—we need not write this as a partial derivative with fields held constant [cf. (2)]. The term $\sum_{\phi \in \{f\}} \gamma^{(\phi)}$ explicitly takes account of

the anomalous dimensions of the fields which suffer field strength renormalization. The field ϕ belongs to the set of fields $\{f\}$ and the notation $\gamma^{(\phi)}$ just stands for the anomalous dimension of the field ϕ (which, we recall, is zero for the gauge field, as a consequence of the manifest gauge invariance).

The lobes on the right-hand side of the flow equation are vertex coefficient functions of S and Σ_g . These lobes are joined together by the ERG kernels, $\text{---}\bullet\text{---}$, which are covariantized, as appropriate. In QED it is necessary to covariantize only the kernels of the fermions and their PV partners, meaning that these kernels can be decorated by gauge fields. The rule for decorating the diagrams on the right-hand side is simple: the set of fields, $\{f\}$, are distributed in all allowed, independent ways between the component objects of each diagram. For the details, the reader is referred to [6].

The understanding and efficient application of the diagrammatic flow equation has been tremendously enhanced through a diagrammatic calculus, proposed in [13], refined in [6, 11, 18–20, 22–25] and completed in [21]. The central ingredient of this calculus is the ‘effective propagator relation’. The recent understanding of this relation [8, 21] is as follows. Starting with the kernels, $\hat{\Delta}$, introduce the integrated kernels, Δ , such that

$$-\Lambda \frac{d}{d\Lambda} \Delta \equiv \hat{\Delta}.$$

The integrated kernels are what we refer to as the effective propagators. Next, define a set of two-point vertices, Δ^{-1} , that are essentially the inverses of the effective propagators. Indeed, in the fermion and PV sectors, these vertices are precisely the inverses of the corresponding effective propagators, but in the gauge sector things are more subtle. As the name suggests, effective propagators are somewhat like usual propagators. In the fermion and PV sectors, they can be taken to be precisely UV regularized propagators [6]. In the gauge sector, however, it is clear that we cannot interpret the integrated kernel simply as a regularized propagator, since we have not fixed the gauge and so cannot define a propagator in the usual sense! Nevertheless, there is nothing to stop us from defining ERG kernels and integrating them with respect to the effective scale. Now, when we come to contract the gauge sector effective propagator into the $(\Delta^{-1})_{\mu\nu}(p)$ vertex, we should get the identity *plus a remainder term*, where this remainder is forced upon us by gauge invariance. (This is all explained more fully in [6, 13, 21].) Specifically, the gauge sector effective propagator is the inverse of the appropriate two-point vertex, in the transverse space:

$$(\Delta^{-1})_{\mu\nu}(p) \Delta^{AA}(p) = \delta_{\mu\nu} - \frac{p_\mu p_\nu}{p^2}. \quad (4)$$

To give a specific example, let us introduce the UV cutoff function, $c(p)$, which satisfies $c(0) = 1$ and dies off sufficiently fast as $p^2/\Lambda^2 \rightarrow \infty$. We could now choose to

identify $(\Delta^{-1})_{\mu\nu}(p)$ with the regularized classical two-point vertex, $c^{-1}(p) \square_{\mu\nu}(p)$ and take $\Delta^{AA}(p) = c(p)/p^2$, which clearly satisfies (4) (we have defined $\square_{\mu\nu}(p) \equiv p^2 \delta_{\mu\nu} - p_\mu p_\nu$).

The reason that the effective propagator relation is so useful is that it allows the simplification of a certain class of diagrams, which then allows the cancellation of other diagrams generated in a typical calculation (see, particularly, [22]). In the fermion and regulator sectors, this is the end of the story. In the gauge sector, we are left over with the remainders. However, it turns out that these can be processed diagrammatically, using the Ward identities [6, 22], and the whole procedure of applying the effective propagator relation and cancelling terms can be iterated. As we shall see shortly, one result of these cancellations is that the β -function possesses no explicit dependence on either the seed action or the details of the covariantization of the cutoff. Looking at figure 1, this is really rather remarkable. Given all these cancellations, what is it that the β -function depends on? The answer is simply the exact n -point vertices, with all instances of Δ^{-1} having been extracted, joined together by effective propagators. Indeed, since instances of Δ^{-1} are removed via application of the effective propagator relation, it is useful to define reduced vertices according to:

$$\left[\text{SR} \right]^{\{f\}} \equiv \left[\text{S} - \frac{1}{g^2} \text{S} \Delta^{-1} \delta_{2,n_f} \right]^{\{f\}}, \quad (5)$$

where n_f is the number of fields in the set $\{f\}$. If we chose to identify the Δ^{-1} vertices with the canonical classical, two-point vertices, then the reduced vertices are simply the vertices of the interaction part of the Wilsonian effective action.

Our aim, now, is to use the diagrammatic form of the flow equation to compute the flow of a special combination of diagrams. Following [8, 21–23], consider the following diagrammatic expression, which basically constitutes all connected diagrams, possessing two external gauge fields, and built from *exact* n -point vertices:

$$\mathcal{D}_{\mu\nu}(p) \equiv \sum_{s=0}^{\infty} \sum_{j=1}^{s+1} \Upsilon_{s,j} g^{2s} \left[\left[\text{SR} \right]^j \right]^{\Delta^s A_\mu(p) A_\nu(-p)} \quad (6)$$

where, for non-negative integers a and b ,

$$\Upsilon_{a,b} \equiv \frac{(-1)^{b+1}}{a!b!} \left(\frac{1}{2} \right)^a. \quad (7)$$

We understand the notation of (6) as follows. The right-hand side stands for all independent, connected diagrams which can be created from j reduced Wilsonian effective action vertices, s internal lines (i.e. effective propagators) and which are decorated by two external gauge fields, carrying momenta p and $-p$ into the vertex. (It is the constraint of connectedness which restricts the sum over j .) The combinatorics for generating fully fleshed out

diagrams is simple and intuitive and is described in [8] (see also [22, 24]). To gain a feeling for the structure of (6), the first few terms represented by the right-hand side are shown in figure 2.

FIG. 2: The first few terms contributing to $\mathcal{D}_{\mu\nu}(p)$ (Lorentz indices and momenta on the external lines have been suppressed). The flavours of the internal fields are essentially summed over; for the precise statement of the Feynman rules, see [6]. Since reduction of the vertices only affects two-point vertices, we have removed the superscript ‘R’ from all vertices with more than two legs.

Note that the diagrams of figure 2 have certain similarities to standard Feynman diagrams; indeed, were we to shrink the lobes to points, they would look the same. However, despite this similarity, our diagrams are related to the ERG flows of vertices of exact Wilsonian effective action and not (directly) to perturbative scattering amplitudes. Whilst physics can most certainly be extracted from the Wilsonian effective action vertices, in the current case this must be done in a manifestly gauge invariant way [23].

Using the diagrammatic calculus it is straightforward, but somewhat tedious, to demonstrate that:

$$\frac{2\beta}{g^3} \square_{\mu\nu}(p) + \mathcal{O}(p^4) = \Lambda \frac{d}{d\Lambda} \mathcal{D}_{\mu\nu}(p), \quad (8)$$

where

$$\beta \equiv \Lambda \frac{dg}{d\Lambda}. \quad (9)$$

(An explicit demonstration of many of the steps pertinent to this calculation are shown in the simpler case of scalar field theory in [8]. See also [22].) Note that, as claimed earlier, the β -function has been written in a form where there is no explicit dependence on either the seed action or the details of the covariantization of the cutoff.

At this stage, we would do well to pause and carefully assess what kind of nonperturbative conclusions we can reliably draw from (8). The first point to make is that the entire diagrammatic approach relies on a weak field expansion, which has its drawbacks. For example, searches for nonperturbative fixed points, using such a scheme, rely on truncating the infinite tower of coupled equations for the exact n -point vertices, and this is known to give

bad results [26] (but see also [27]). However, the situation is much better in the current case, as we now argue. First, we never perform any truncations and will instead draw conclusions from general properties of the full function $\mathcal{D}_{\mu\nu}(p)$, which we emphasise depends on the *exact* n -point vertices, no perturbative expansions having been performed. Secondly, we can always consider this function in the weak coupling regime. This does not mean to say that we wish to do perturbation theory, throwing away all nonperturbative contributions. Rather, we simply want to consider $\mathcal{D}_{\mu\nu}(p)$ in a regime where its diagrammatic expansion can be ordered with a small parameter and could, at least in principle, be exactly resummed since we have not thrown any contributions away.

Thus, our understanding of $\mathcal{D}_{\mu\nu}(p)$ is as follows. Formally, it is given by its diagrammatic expansion (6) at all scales. More rigorously, this diagrammatic expansion should be evaluated in the regime where the coupling is small, and resummed.

With these points in mind, we will now show that there cannot, in fact, be nonperturbative contributions to (8), implying that the perturbative expansion of the β -function can actually be resummed, by itself. To do this, we re-express (8) as:

$$\frac{2\beta}{g^3} + \mathcal{O}(p^2) = \frac{\Lambda \partial_\Lambda \mathcal{D}'(p)}{1 - g^3/2\partial_g \mathcal{D}'(p)}, \quad (10)$$

where $\mathcal{D}'(p) \square_{\mu\nu}(p) \equiv \mathcal{D}_{\mu\nu}(p)$ and the partial derivative with respect to Λ is performed at constant g . We now make the following observation: loop integrals in the diagrams comprising $\mathcal{D}'(p)$ can acquire factors of $\ln p^2/\Lambda^2$, arising from IR divergences in the limit $p \rightarrow 0$. This is clear from analysing e.g. the third diagram in figure 2, for which the component which goes like

$$\int d^4k \frac{1}{k} \frac{1}{p+k}$$

in the IR (see [6] for the details) produces the desired behaviour after we act on the full diagram with the Λ -derivative and pull out $\square_{\mu\nu}(p)$. [On dimensional grounds, we see that the integrand $\sim p^2/k^4$, at $\mathcal{O}(p^2)$.] It is important to note that the apparent UV divergence in $\ln p^2/\Lambda^2$ is an artefact of us having Taylor expanded vertices and cutoff functions in the external momentum (as it must be: by construction, everything is UV regularized). Indeed, as mentioned earlier, all vertices have a derivative expansion, as do the cutoff functions. The only non-polynomial dependence of $\mathcal{D}'(p)$ on the external momentum is generated by certain loop integrals in the IR.

Furthermore, whilst individual diagrams contributing to $\mathcal{D}'(p) + \mathcal{O}(p^2)$ can diverge as a logarithm to a power (which is at most equal to the number of loops) as $p \rightarrow 0$, all divergences must cancel out between the numerator and denominator of (10) since the $\mathcal{O}(p^0)$ contribution to the left-hand side of (10) has no additional, non-polynomial dependence on p . Consequently, for functions

F_1 , F_2 and G , it must be that we can write:

$$\frac{2\beta}{g^3} + \mathcal{O}(p^2) = \frac{F_1(g^2)G(g^2, \ln p^2/\Lambda^2)}{F_2(g^2)G(g^2, \ln p^2/\Lambda^2)} = \frac{F_1(g^2)}{F_2(g^2)}. \quad (11)$$

So, to begin with, let us consider perturbative contributions in (10). Let us suppose that, at order g^{2n} , the strongest IR divergence carried by $\mathcal{D}'(p)$ goes like

$$g^{2n} \ln^m p^2/\Lambda^2. \quad (12)$$

In the numerator, the Λ -derivative reduces this divergence to one of the form

$$g^{2n} \ln^{m-1} p^2/\Lambda^2 \quad (13)$$

whereas, in the denominator, a contributions of the form

$$g^{2(n+1)} \ln^m p^2/\Lambda^2 \quad (14)$$

is produced. Thus, we have found that terms of the form (12) provide a divergent contribution to the denominator which does not seem to exist in the numerator. Of course, there is no real problem here: all we need to do is consider diagrams with an extra loop. In such diagrams there are contributions of the form (12) but with $n \rightarrow n+1$ and $m \rightarrow m+1$. Terms like this in the numerator are, after differentiation with respect to Λ , of precisely the right form to cancel denominator contributions of the type (14). This is explicitly borne out in perturbative calculations [18, 19].

But now consider a contribution of the type

$$g^{2n} e^{-a/g^2} \ln^m p^2/\Lambda^2, \quad (15)$$

where again we assume that, for our choice of n , there is no stronger IR divergence. In the numerator this contributes terms of the form

$$g^{2n} e^{-a/g^2} \ln^{m-1} p^2/\Lambda^2 \quad (16)$$

and in the denominator it yields terms of the form

$$g^{2n} e^{-a/g^2} \ln^m p^2/\Lambda^2 + \dots, \quad (17)$$

where the ellipsis denotes terms higher order in g^2 . Crucially, (16) and (17) are *the same order* in g^2 . Since, by assumption, there are no terms in $\mathcal{D}'(p)$ which are of order $g^{2n} e^{-a/g^2}$ but which have a stronger IR divergence than (15), there is no way that the denominator contribution (17) can ever be cancelled. From (11), we therefore conclude that terms of the type (15) must be absent from (11), unless $m=0$. But it is easy to see that $m=0$ terms can appear only in $G(g^2, \ln p^2/\Lambda^2)$ and not in $F_1(g^2)$ or $F_2(g^2)$: for if this condition is violated, then we necessarily produce contributions of the form (15), when we expand out $F_1(g^2)G(g^2, \ln p^2/\Lambda^2)$. In conclusion, the only contributions to the β -function of the form (15) that are allowed—namely those with $m=0$ —cancel out!

It is now straightforward to generalize this argument to show that only the perturbative contributions to the β -function survive. First, we note that the above argument is not affected if we consider terms which include e^{-b/g^4} , e^{-c/g^6} etc., or products of such terms. Secondly, we can allow additional functions of g to come along for the ride, so long as they do not spoil the requirement that the ERG trajectory sinks into the Gaussian fixed point as $\Lambda \rightarrow 0$.

Thus, we have demonstrated that the β -function is free of nonperturbative power corrections and, therefore, must be resumable, at least in the massless theory. In the presence of a fermion mass, μ , there would be no good reason to exclude surviving contributions to the β -function which go like

$$\frac{\mu^2}{\Lambda^2} e^{-a/g^2},$$

since the mass now regularizes terms which previously diverged as $p \rightarrow 0$. Note, though, that as emphasised in [7], the presence of exponentially small terms does not, by itself, necessarily imply that perturbation theory cannot be resummed (though it is suggestive). In other words, it is at least possible that a (non-analytic) function comprises a resumable perturbative series plus additional exponentially small terms. Of course, were this scenario to be realized in the massive case, one would certainly have to provide an argument as to why the perturbative series was free of renormalons.

Returning to the massless theory, the resumability of the β -function is valid in the infinite number of schemes implicit to our approach; these different schemes corresponding to all the legal choices of the seed action and covariantization of the cutoff. There is no reason to expect that this conclusion is true for unrelated schemes, such as \overline{MS} . It is important to add that no expression as neat as (8) exists for the flow of the other couplings or for the anomalous dimension, γ (see [6, 8, 22] for the tools necessary to compute these expressions). Consequently, there is no way to argue that the perturbative series for these functions, also, are resumable. Indeed, we expect precisely the opposite to be true, since we do not expect self-similar trajectories to exist within the critical surface of the Gaussian fixed point. Nevertheless, it would doubtless be interesting to compute the β -function to some high order in perturbation theory and resum it, not least from the point of view of understanding the fate of the Landau pole in the massless theory. Perhaps more interesting still would be to try to get some handle on what happens in the massive case, particularly given the work already done on ERG flows in QED [28].

Finally, we should note that one can attempt to repeat the above analysis for other field theories. In QCD, the expression for the β -function possesses additional terms, which can spoil the above arguments (the basic structure is apparent at the perturbative level [22]). Nevertheless, for this to happen, there must be some delicate relationships between the various terms contributing to the β -function, which merits further investigation [9]. In scalar

field theory the β -function has a sufficiently complicated form to spoil the above arguments, as expected [1, 29]. The most interesting case to look at, as mentioned earlier, is the Wess-Zumino model, where the above resummability argument seems to apply to the anomalous dimension (equivalently the β -function if we induce a flow of the superpotential by using the field strength renormalization to rescale the field). This is exciting because, unlike in QED, there are arguments to suggest that resummability of the anomalous dimension implies resummability of the

entire perturbative expansion of the theory. If true, this would be highly suggestive of the existence of a UV fixed point [9].

Acknowledgments

I acknowledge IRCSET for financial support. I would like to thank Denjoe O'Connor for useful discussions.

-
- [1] T. R. Morris, “Elements of the continuous renormalization group,” *Prog. Theor. Phys.* **131** (1998) 395, [hep-th/9802039](#).
- [2] C. Bagnuls and C. Bervillier, “Exact renormalization group equations: An introductory review,” *Phys. Rept.* **348** (2001) 91, [hep-th/0002034](#).
- [3] K. Wilson and J. Kogut, “The Renormalization group and the epsilon expansion,” *Phys. Rept.* **12** (1974) 75.
- [4] F. J. Wegner and A. Houghton, “Renormalization group equation for critical phenomena,” *Phys. Rev. A* **8** (1973) 401.
- [5] P. Hasenfratz and F. Niedermayer, “Perfect Lattice Action For Asymptotically Free Theories,” *Nucl. Phys. B* **414** (1994) 785, [hep-lat/9308004](#).
- [6] S. Arnone, T. R. Morris, and O. J. Rosten, “Manifestly gauge invariant QED,” *JHEP* **0510** (2005) 115, [hep-th/0505169](#).
- [7] M. Beneke, “Renormalons,” *Phys. Rept.* **317** (1999) 1, [hep-th/9807443](#).
- [8] O. J. Rosten, “Sensitivity of Nonrenormalizable Trajectories to the Bare Scale,” [arXiv:0710.3658 \[hep-th\]](#).
- [9] O. J. Rosten, “Work in Progress.”
- [10] T. R. Morris and J. L. Latorre, “Exact scheme independence,” *JHEP* **0011** (2000) 004, [hep-th/0008123](#).
- [11] S. Arnone, T. R. Morris, and O. J. Rosten, “A Generalised manifestly gauge invariant exact renormalisation group for SU(N) Yang-Mills,” *Eur. Phys. J. C* **50** (2007) 467, [hep-th/0507154](#).
- [12] T. R. Morris, “A gauge invariant exact renormalization group. I,” *Nucl. Phys. B* **573** (2000) 97, [hep-th/9910058](#).
- [13] S. Arnone, A. Gatti, and T. R. Morris, “A proposal for a manifestly gauge invariant and universal calculus in Yang-Mills theory,” *Phys. Rev. D* **67** (2003) 085004, [hep-th/0209162](#).
- [14] T. R. Morris and O. J. Rosten, “Manifestly gauge invariant QCD,” *J. Phys. A* **39** (2006) 11657, [hep-th/0606189](#).
- [15] S. Arnone, T. R. Morris, and O. J. Rosten, “Manifestly gauge invariant exact renormalization group,” *Fields Institute Communications* **50** (2007) 1, [hep-th/0606181](#).
- [16] J. Polchinski, “Renormalization And Effective Lagrangians,” *Nucl. Phys. B* **231** (1984) 269.
- [17] L. P. Kadanoff, “Scaling laws for Ising models near T(c),” *Physics* **2** (1966) 263.
- [18] T. R. Morris and O. J. Rosten, “A manifestly gauge invariant, continuum calculation of the SU(N) Yang-Mills two-loop beta function,” *Phys. Rev. D* **73** (2006) 065003, [hep-th/0508026](#).
- [19] S. Arnone, A. Gatti, T. R. Morris, and O. J. Rosten, “Exact scheme independence at two loops,” *Phys. Rev. D* **69** (2004) 065009, [hep-th/0309242](#).
- [20] O. J. Rosten, *The manifestly gauge invariant exact renormalisation group*. PhD thesis, Southampton U., 2005. [hep-th/0506162](#).
- [21] O. J. Rosten, “Universality from very general nonperturbative flow equations in QCD,” *Phys. Lett. B* **645** (466) 2007, [hep-th/0611323](#).
- [22] O. J. Rosten, “A manifestly gauge invariant and universal calculus for SU(N) Yang-Mills,” *Int. J. Mod. Phys. A* **21** (2006) 4627, [hep-th/0602229](#).
- [23] O. J. Rosten, “General computations without fixing the gauge,” *Phys. Rev. D* **74** (2006) 125006, [hep-th/0604183](#).
- [24] O. J. Rosten, “Scheme independence to all loops,” *J. Phys. A* **39** (2006) 8699, [hep-th/0507166](#).
- [25] O. J. Rosten, “A primer for manifestly gauge invariant computations in SU(N) Yang-Mills,” *J. Phys. A* **39** (2006) 8699, [hep-th/0507166](#).
- [26] T. R. Morris, “On truncations of the exact renormalization group,” *Phys. Lett. B* **334** (1994) 355, [hep-th/9405190](#).
- [27] K. I. Aoki, K. Morikawa, W. Souma, J. I. Sumi, and H. Terao, “Rapidly converging truncation scheme of the exact renormalization group,” *Prog. Theor. Phys.* **99** (1998) 451, [hep-th/9803056](#).
- [28] H. Gies and J. Jaeckel, “Renormalization flow of QED,” *Phys. Rev. Lett.* **93** (2004) 110405, [hep-ph/0405183](#).
- [29] T. R. Morris, “The Exact renormalization group and approximate solutions,” *Int. J. Mod. Phys. A* **9** (1994) 2411, [hep-ph/9308265](#).